WHEAT (INCLUDING DURUM), BARLEY, OATS - SUSCEPTIBLE BROADLEAF WEED SEEDLINGS LESS THAN 4 INCHES TALL

General Information

GNERAL INFORMATION

Application Precautions

- Avoid application where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift.

- Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, or drainage. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the intended rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop or crop listed in the table below for which the rotational interval has clearly been met.

Application Restrictions

- Do not apply FULL DECK directly to, or allow spray drift to come in contact with broadleaf crops or other susceptible broadleaf plants, including, but not limited to, alfalfa, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season.

- Do not contaminate irrigation ditches or water used for domestic purposes.

- Chemigation: Do not apply this product through any type of irrigation system.
- Do not transfer livestock from treated grazing areas (or feeding of treated hay) to

sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Avoiding Injury to Non-Target Plants: This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Do not apply FULL DECK herbicide directly to, or allow spray drift to come in contact with broad leaf crops, including, but not limited to alfalfa, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See guidance in section entitled "Crop Rotation Intervals".)

Residues in Plants or Manure: Do not use plant residues, including hay or straw from treated areas, or manure or bedding straw from animals that have grazed or consumed forage from treated areas, for composting or mulching, where susceptible plants may be grown the following season. Do not spread manure from animals that have grazed or consumed forage or hay from treated areas on land used for growing susceptible broadleaf crops. To promote herbicidal decomposition, plant residues should be evenly incorporated or burned. Breakdown of clopyralid in crop residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.

Avoid Movement of Treated Soil: Avoid conditions under which soil from treated areas may be moved or blown to areas containing susceptible plants. Wind-blown dust containing clopyralid may produce visible symptoms, such as epinasty (downward curving or twisting of leaf petioles or stems) when deposited on susceptible plants; however, serious injury is unlikely. To minimize potential movement of clopyralid on wind-blown dust, avoid treatment of powdery dry or light sandy soils until soil has been settled by rainfall or irrigation or irrigate shortly after application.

Precautions for Avoiding Spray Drift: Spray drift, even very small quantities of the spray that may not be visible, may severely injure susceptible crops whether dormant or actively growing. When applying FULL DECK, use low-pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets. Under adverse weather conditions, fine spray droplets that do not settle rapidly onto target vegetation may be carried a considerable distance from the treatment area. A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Applications: To minimize spray drift, apply FULL DECK in a total spray volume of 8 or more gallons per acre using spray equipment designed to produce large-droplet, low pressure sprays. Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Spot treatments should be applied only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.

Aerial Application: To minimize spray drift, apply FULL DECK in a total spray volume of 3 or more gallons per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high potential for temperature inversion. Spray drift from aerial application can be minimized by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 3/4 the rotor or wing span of the aircraft. Spray pattern and droplet size distribution can be evaluated by applying sprays containing a water soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Do not apply under conditions of a low level air temperature inversion. A temperature inversion is characterized by little or no wind and lower air temperature near the ground than at higher levels. The behavior of smoke generated by an aircraft mounted device or continuous smoke column released at or near site of application will indicate the direction and velocity of air movement. A temperature inversion is indicated by layering of smoke at some level above the ground and little or no lateral movement.

Perennial weeds: FULL DECK will control the initial top growth and inhibit regrowth

during the season of application (season long control). At higher use rates shown on the label, FULL DECK may cause a reduction in shoot regrowth in the season following application; however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

Management of Kochia Biotypes: Research has suggested that many biotypes of kochia can occur within a single field. While kochia biotypes can vary in their susceptibility to FULL DECK, all will be suppressed or controlled by the 1.0 pint per acre labeled rate. Application of FULL DECK at rates below the 1.0 pint per acre rate can result in a shift to more tolerant biotypes within a field.

Best Resistance Management Practices: Extensive populations of dicamba tolerant kochia have been identified in certain small grain production regions (such as Chouteau, Fergus, Liberty, Toole, and Treasure counties in the state of Montana). For optimal control of dicamba tolerant kochia in these counties, FULL DECK is recommended at a minimum rate of 1.0 pints per acre. In addition, use of FULL DECK should be rotated with products that do not contain dicamba to minimize selection pressure. Use of these practices will preserve the utility of FULL DECK for control of dicamba tolerant kochia biotypes.

Spot Treatments: To prevent misapplication, it is recommended that spot treatments be applied only with a calibrated boom or with hand sprayers according to directions provided below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq ft. Mix the amount of FULL DECK (fl oz or ml) corresponding to the desired broadcast rate in 10 or more gallons of spray. To calculate the amount of FULL DECK required for larger areas, multiply the table value (fl oz or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3,500 sq ft, multiply the table value by 3.5 (calc. 3,500 \div 1,000 = 3.5). An area of 1000 sq ft is approximately 10.5 x 10.5 yards (strides) in size.

Limitations, Restrictions, and Exceptions

WHEAT (including Durum), BARLEY, OATS

Application Timing: Apply to actively growing weeds. Extreme growing conditions

such as drought or near freezing temperatures prior to, at, or following application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds that have emerged at the time of application will be controlled. If foliage is wet at the time of application, control may be decreased. Applications of FULL DECK herbicide are rainfast within 6 hours after application. To obtain seasonlong control of perennial weeds such as Canada thistle, apply when the majority of the basal leaves have emerged from the soil up to bud stage. For suppression of volunteer potatoes, apply before potato plants are 6 inches tall. Do not apply FULL DECK after the crop has reached the F9 stage. Do not use if cereal crop is underseeded with a legume.

Effect of Temperature on Herbicidal Activity: Herbicidal activity of FULL DECK is influenced by weather conditions. Optimum activity requires active plant growth. The temperature range for optimum herbicidal activity is 55°F to 75°F. Reduced activity will occur when temperatures are below 45°F or above 85°F. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance.

Restrictions:

Do not allow livestock to graze treated areas or harvest treated forage within 7 days of application.

Do not apply more than 1.7 pints per acre of FULL DECK per growing season.

Preharvest Interval: Do not apply closer than 14 days before cutting of hay or 40 days before harvesting of grain and straw.

Application Rates: Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of susceptible weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rates within the rate range will be needed. Weeds in fallow land or other areas where competition from crops is not present will generally require higher rates for control or suppression.

Maximum MCPA: 0.310 lbs a.e./acre

Weeds Controlled

Artichoke, Jerusalem, Cocklebur, Common, Jimsonweed, Marshelder, Ragweed, common, Ragweed, Giant, Sunflower: For best control, apply up to 5 leaf stage of growth.

Bedstraw (cleavers): For best control, apply in the 1-4 leaf "whorl" stage of growth.

Buckwheat, wild: For best control, apply in the 1– 3 leaf stage of growth, before vining.

Kochia: Includes herbicide tolerant or resistant biotypes. Best control is achieved when weeds are at least 1 inch tall.

Nightshade, black: For best control or suppression, apply at the 2 – 4 leaf stage of growth.

Nightshade, Cutleaf, nightshade, hairy, buffalobur, ladysthumb, Thistle, Canada: For best control or suppression, apply from rosette to bud (pre-flower) stage of growth.

Weeds Suppressed: Suppression is expressed as a reduction in weed competition (reduction population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

Method <u>Broadcast/Foliar Air</u> <u>Broadcast/Foliar Ground</u> <u>Spot treatment</u> Rates field rates 0

• Restricted Entry Interval

12 hours

Timings Postemergence (Weed)