

SPOT OR LAKE MARGIN TREATMENT - COONTAIL, HORNED PONDWEED, ETC.

General Information

PRODUCT INFORMATION

Aquathol K is a liquid concentrate soluble in water which is effective against a broad range of aquatic plants. Dosage rates indicated for the application of Aquathol K are measured in parts per million (ppm) of dipotassium endothall.

DIRECTIONS FOR USE

- For quiescent or slow moving water treatments: Waters treated with Aquathol K may be used for swimming, fishing, and irrigating turf, ornamental plants and crops immediately after treatment with the following exceptions: Do not use the Aquathol K treated water to irrigate the following for 7 days after the treatment: annual nursery or greenhouse crops including hydroponics and newly seeded or transplanted annual crops, newly seeded or transplanted ornamentals, and newly sodded or seeded turf. Do not use treated water for animal consumption within the following periods:

0.5 ppm dipotassium salt – 7 days after application

4.25 ppm dipotassium salt – 14 days after application

5.0 ppm dipotassium salt – 25 days after application

- For flowing water treatments: Waters treated with Aquathol K may be used for swimming, fishing, livestock watering, and irrigating turf, ornamental plants and crops immediately after treatment with the following exceptions: Do not use the Aquathol K treated water to irrigate the following: annual nursery or greenhouse crops including hydroponics and newly seeded or transplanted annual crops, newly seeded or transplanted ornamentals, and newly sodded or seeded turf.

- Phytotoxicity is not expected on plants or crops irrigated with Aquathol K treated water, however, all species and cultivars (varieties) have not been tested.

- Undiluted Aquathol K may be injurious to crops, grass, ornamentals, and other foliage.
- Do not use Aquathol K treated water for chemigation as interactions between Aquathol K and other pesticides and fertilizers are not known.
- Do not use Aquathol K in brackish or saltwater.
- Wash out spray equipment with water after each operation.
- Avoid contact of spray concentrate (product) directly or by drift with non-target plants or crops as injury may result.

HOW TO APPLY:

Aquathol K is a contact herbicide; consequently, apply when target plants are present.

Aquathol K should be sprayed on the water or injected below the water surface. It may be applied as a concentrate or diluted with water depending on the equipment.

In instances where the plant(s) to be controlled is an exposed surface problem (i.e., some of the broad-leaved pond weeds) coverage is important. For best results, apply the concentrate with the least amount of water compatible with the application equipment.

Drinking Water (Potable Water)

Consult with appropriate state or local water authorities before applying this product to public waters. State or local agencies may require permits.

The drinking water (potable water) restrictions on the label are to ensure that consumption of water by the public is allowed only when the concentration of endothall acid in the water is less than the MCL (Maximum Contamination Level) of 0.1 ppm. Applicators should consider the unique characteristics of the treated waters to assure that endothall acid concentrations in potable drinking water do not exceed 0.1 ppm at the time of consumption.

For Lakes, Ponds, and other Quiescent Water Bodies:

- For Aquathol K applications, the drinking water setback distance from functioning

potable water intakes in the treated water body must be greater than or equal to 600 feet.

- Note: Existing potable water intakes that are no longer in use, such as those replaced by a connection to a municipal water system or a potable water well, are not considered to be functioning potable water intakes.

For Irrigation Canals and other Flowing Water Bodies:

- Applicator is responsible to assure that treated water does not enter potable water intakes. For Aquathol K applications, potable water intakes must be closed when treated water is present at the intake. In the event the water intake cannot be closed, treatments must only be made downstream from the intake in order to assure Aquathol K treated water does not enter the potable water system.

Limitations, Restrictions, and Exceptions

QUIESCENT OR SLOW MOVING WATER TREATMENTS: SURFACE OR INJECTED APPLICATIONS

For aquatic plant control in quiescent or slow moving water, Aquathol K recommended use rates can be found in the following chart. Since the active ingredient is water soluble and tends to diffuse from the treated area, select the dosage rate applicable to the area to be treated. Marginal treatments of large bodies of water require higher rates as indicated.

Use higher labeled rates of Aquathol K when making treatments to small areas with an increased potential for rapid dilution or when treating narrow areas such as boat lanes or shoreline treatments where dilution may reduce the exposure of plants to Aquathol K. Use lower labeled rates of Aquathol K for large contiguous treatment blocks or in protected areas such as coves where reduced water movement will not result in rapid dilution of Aquathol K from the target treatment area or when treating entire lakes or ponds.

See label for the charts indicated the quantity of Aquathol K to be applied.

Restrictions: Do not apply more than a total of 5 ppm within a 7-day interval.

Note: There is no Pre-harvest Interval (PHI) for crops irrigated with treated water.

Method

[Spot treatment](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[When target plants are present.](#)