

## **ORNAMENTALS - LANDSCAPE PLANTINGS**

### **General Information**

**DIRECTIONS FOR USE** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not formulate this product into other end-use products.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

PROKOZ ZENITH 0.5 G Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

### **APPLICATION EQUIPMENT**

Apply PROKOZ ZENITH 0.5 G Insecticide uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply PROKOZ ZENITH 0.5 G Insecticide. Do not use spreaders that would apply the material in concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

### **RESTRICTIONS**

Do not apply PROKOZ ZENITH 0.5 G Insecticide to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant. Do not apply more than 80 lb (0.4 lb of active ingredient) per acre per year.

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must

be observed.

## Limitations, Restrictions, and Exceptions

### APPLICATION TO ORNAMENTALS

PROKOZ ZENITH 0.5 G Insecticide can be used as a broadcast application on ornamentals in commercial and residential landscapes. PROKOZ ZENITH 0.5 G Insecticide is a systemic product and will be translocated upward within the plant system. Optimum control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation, or rainfall to move the active ingredient into the soil. Make application prior to anticipated pest infestation to achieve optimum levels of control.

Restrictions: Not for use in commercial greenhouses, nurseries, on sod farms or on grass grown for seed. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens, or parks, or on golf courses or lawns and grounds.

Do not apply more than 80 lb (0.4 lb of active ingredient) per acre per year.

Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.

NOTE: For optimum control mechanical incorporation, irrigation or rainfall should occur within 24 hours after application to move the active ingredient into soil. Avoid runoff or puddling of irrigation water following application.

Refer to label sections on application equipment and spreader settings.

Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.

### Method

#### [Broadcast](#)

### Rates

#### [field\\_rates 0](#)

- 

### Timings

#### [N. A.](#)