

POA ANNUA CONTROL IN PERENNIAL RYEGRASS OVERSEEDING (ALABAMA, LOUISIANA, ETC.)

General Information

GENERAL INFORMATION

WHERE TO USE

PRODIAMINE 65 WDG is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turfgrass (excluding golf course putting greens), lawns, and sod nurseries
- Container, field-grown, and landscape ornamentals
- Conifer and hardwood seedling nurseries
- Established perennial and wildflower plantings
- Noncrop areas including managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways, and equipment yards)
- Facilities including substations, tank farms, pumping stations, parking and storage areas, and ungrazed fence rows
- Christmas tree farms

Drainage ditches in California and Arizona only

HOW PRODIAMINE 65 WDG WORKS

PRODIAMINE 65 WDG controls susceptible weeds by preventing growth and development of newly germinated weed seeds. Weed control is most effective when PRODIAMINE 65 WDG is activated by at least 0.5 inch of rainfall or irrigation or shallow incorporation (1- 2 inches) before weed seeds germinate and within 14 days following application.

USE PRECAUTIONS

- Do not graze or feed livestock forage cut from areas treated with Quali-Pro Prodiamine 65 WDG.
- Do not apply Quali-Pro Prodiamine 65 WDG to plants that will be consumed for food use.
- Follow all applicable directions, restrictions, and precautions on the labels of EPA-registered tank mix partners.
- Do not blend Quali-Pro Prodiamine 65 WDG onto dry fertilizer or any other granular material.
- Chemigation Statement: do not apply this product through any type of irrigation system unless instructed otherwise in the label.
- Do not apply aerially.
- Do not apply to golf course putting greens.

Limitations, Restrictions, and Exceptions

WHEN TO APPLY QUALI-PRO PRODIAMINE 65 WDG AFTER OVERSEEDING TURF

Injury to desirable seedlings is likely if Quali-Pro Prodiamine 65 WDG is applied before the secondary roots of seedlings are in the second inch of soil (not thatch plus soil). To reduce the potential to injure overseeded turf, wait 60 days after seeding or until after the second mowing, whichever is longer, before applying Quali-Pro Prodiamine 65 WDG.

When to Overseed After Application (All States)* Quali-Pro Prodiamine 65 WDG will inhibit the development of turfgrass species overseeded too soon after application. Follow rates and intervals in the table for best overseeding/reseeding results.

*Note: See exceptions for "Poa Annua Control in Established Bermudagrass Overseeded with Perennial Ryegrass" in the label.

EXPECTED CONTROL

- 70% or greater

USE PRECAUTIONS

- Some seedling mortality and temporary reduction in root growth of new seedlings may occur.
- To reduce the potential for seedling mortality, maintain a moist seedbed with light, frequent irrigation.
- To maximize seedling establishment, use lower rate and/or the maximum time interval before overseeding. To maximize *Poa annua* control, use higher rate and shorter time interval before overseeding.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If this product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[6 to 8 weeks before ryegrass overseeding](#)