

# **SOUTHERN PINE SEEDBEDS**

## General Information

### GENERAL INFORMATION

#### WHERE TO USE

PRODIAMINE 65 WDG is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turfgrass (excluding golf course putting greens), lawns, and sod nurseries
- Container, field-grown, and landscape ornamentals
- Conifer and hardwood seedling nurseries
- Established perennial and wildflower plantings
- Noncrop areas including managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways, and equipment yards)
- Facilities including substations, tank farms, pumping stations, parking and storage areas, and ungrazed fence rows
- Christmas tree farms

Drainage ditches in California and Arizona only

#### HOW PRODIAMINE 65 WDG WORKS

PRODIAMINE 65 WDG controls susceptible weeds by preventing growth and development of newly germinated weed seeds. Weed control is most effective when PRODIAMINE 65 WDG is activated by at least 0.5 inch of rainfall or irrigation or shallow incorporation (1- 2 inches) before weed seeds germinate and within 14 days following application.

#### USE PRECAUTIONS

- Do not graze or feed livestock forage cut from areas treated with Quali-Pro Prodiamine 65 WDG.
- Do not apply Quali-Pro Prodiamine 65 WDG to plants that will be consumed for food use.
- Follow all applicable directions, restrictions, and precautions on the labels of EPA-registered tank mix partners.
- Do not blend Quali-Pro Prodiamine 65 WDG onto dry fertilizer or any other granular material.
- Chemigation Statement: do not apply this product through any type of irrigation system unless instructed otherwise in the label.
- Do not apply aerially.
- Do not apply to golf course putting greens.

#### Limitations, Restrictions, and Exceptions

#### CONIFER AND HARDWOOD SEEDLING NURSERIES (NONORNAMENTAL, FORESTRY USE ONLY)-VEGETATION MANAGEMENT-QUALI-PRO PRODIAMINE 65 WDG

1. Provides residual preemergence weed control in conifer and hardwood seedling nurseries.
2. Provides the most effective weed control when the product is activated in the soil by 0.5 inch of irrigation or rainfall before weed seeds germinate and within 14 days after application.
3. Should be applied to conifer and hardwood seedling nurseries any time after the soil has settled around newly transplanted seedlings and liners.

#### COMMENTS/INSTRUCTIONS

To assist in the establishment of Southern pine seedbeds, apply this product preemergence just after seeding pines.

Application after emergence of pine seedlings should not occur until 3 weeks after most seedlings have shed their seedcoat.

Mix this product with clean water and broadcast spray at 20 to 40 psi in a minimum of 20 gals. of water per treated area.

After application, sprinkler irrigate beds with approximately 1/2 inch of water.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

12 hours

Exception: If this product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Just after seeding and/or a minimum of 3 weeks after most seedlings have shed their seedcoat.](#)