

## **HIGHLY TOLERANT SPECIES - BROADLEAF WEEDS (SINGLE APPLICATION)**

### General Information

### DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### PRODUCT INFORMATION

Quincept combines Quinclorac, 2,4-D and Dicamba to provide broad-spectrum control of grassy and broadleaf weeds.

- Postemergent control of a comprehensive list of both grassy and broadleaf weeds.
- Excellent clean-up product for areas that did not receive a preemergent grass herbicide application in the spring.
- Controls newly germinated 1 to 2-leaf crabgrass, to 1-tiller crabgrass and when crabgrass has matured to 5-tillers or greater.
- Quinclorac contributes grassy weed control and is absorbed by foliage and roots and translocated throughout the plant.

Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death.

### TIMING

Apply Quincept to actively growing weeds as a postemergence broadcast or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds. Under certain conditions, application of Quincept made to annual grasses at 2 to 4-tiller may not provide complete control.

## NEWLY SEEDED AREAS:

The application of Quincept to grass seedlings is not recommended until after the third mowing.

## NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS:

The application of Quincept to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky Bluegrass, Perennial Ryegrass and Fine Fescues.

## SEEDING:

Delay applications for 4 weeks after seeding and emergence of turf species listed on the label.

## MOWING:

Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

## SUSCEPTIBLE □(DO NOT USE ON THESE GRASS SPECIES)

Bahiagrass, Bentgrass Colonial, Bentgrass Seaside, Buffalograss, Carpetgrass, Centipedegrass, Dichondra, Paspalum Seashore and St. Augustinegrass.

## Limitations, Restrictions, and Exceptions

## USE SITES

Quincept may be applied postemergence to residential and non-residential turfgrasses. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms.

- Warm season grasses such as Common Bermuda and Zoysia may see temporary discoloration. Use reduced rates if grass is stressed from heat or drought. Do not

apply during growth stages from dormancy to green-up and from green-up to dormancy.

- The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.
- Not for use on sod farms in Arizona. For Use in New York by Spot Treatment Only.

#### HIGHLY TOLERANT SPECIES

- Do not exceed 1.5 pounds a.i. Quinclorac per acre per year. Sequential application(s) should be timed 21 days apart.
- Broadleaf weed control may be reduced due to sequential applications. Do NOT use on golf course greens, tees and collars.
- Do NOT use on lawns with desirable clovers or legumes or on ornamentals.

#### APPLICATION GUIDELINES

- Ensure that spray volumes are adequate to completely cover weeds, especially when dense weed infestations make it difficult to completely cover the foliage of target weeds.
- Early applications of this product will not control weeds germinating later in the season so a second application may be necessary.
- If turf injury is observed after the first application, delay the second application until turf has recovered.
- Dual Applications: A single application should control less mature, light weed populations; however, a second application may be required to control heavy, mature broadleaf weed infestations. Under certain conditions, annual grasses in the 2 to 4-tiller stages may require a sequential application. If a second application is required, allow 14 to 21 days between applications.
- For control of woody plants, apply to both stems and foliage any time from the

time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

## SINGLE APPLICATION

### SPRAY VOLUME

- 20 to 300 gal/A or 0.5 to 6.9 gal/1,000 sq. ft.

### MAXIMUM APPLICATIONS

- 2 per year

### Method

[Spray](#)

### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

### Restricted Entry Interval

48 hours

### Timings

[Postemergence \(Weed\)](#)

[when weeds are actively growing and in early stages of growth.](#)