

RATS

General Information

DIRECTIONS FOR USE

Do Not Use in Homes or Other Residential Settings. Do Not Use Outdoors.

USE RESTRICTIONS

For indoor use only. This product is for control of Norway rats, roof rats and house mice only inside of industrial, commercial, agricultural, and public buildings, and similar man-made structures. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA-inspected facilities, this product must be applied in tamper-resistant bait stations. Do not use in sewers.

Do not sell this product in individual containers holding less than 13.44 fl. oz of bait concentrate.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Remove as much alternative food as possible.

MIXING DIRECTIONS: Wearing waterproof gloves, thoroughly mix contents of one 1.68 fl. oz. pouch in 1 quart (32 fluid ounces) of water.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS:

RATS: Provide a minimum of 1 pint (16 fluid ounces) of LIQUA-TOX II diluted product in each dispenser (suitably-equipped bait station, chick fount, or other suitable device). If infested area is large, or if rat population appears to be high, use dispensers at several locations. Space dispensers and refill them as needed to ensure that bait is exposed in all infested areas where it can be used without

putting nontarget organisms at risk.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

Method

[Bait](#)

Timings

[N.A.](#)