

IN AND AROUND BUILDINGS, STRUCTURES, ETC. - FLYING PEST

General Information

PRO FACTS: Ready-to-use, Low-odor, Non-staining

TEMPO 1% DUST Insecticide Ready To Use: Contains 1% cyfluthrin

CONTROLS: Ants, cockroaches, and other crawling pests, Bees and wasps, Pantry and stored product pests, Wood-infesting pests.

COVERAGE: Treats up to 2,500 square feet

WHERE TO APPLY: In and around buildings and structures

HOW TO APPLY: Directly in voids, cracks and crevices. For bees, wasps and ant mounds, see directions inside. Use appropriate dusting or injection equipment.

RE-APPLY: Every 7 to 10-days, if needed

PEST

- For residual pest control in and around buildings, structures, non-food/feed areas of food/feed handling establishments and modes of transport. This includes houses, hotels, motels and other non-food areas of stores, warehouses, hospitals, food processing facilities, and modes of transport.

- The purpose of such applications of TEMPO 1% Dust for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment, or foundation treatment, but merely a supplement. The product is not recommended as sole protection against termites. For active indoor termite infestations get a professional inspection.

WHERE TO APPLY

- Apply where pests have been seen or found or can find shelter.

- In living areas, avoid deposits on exposed surfaces and avoid introducing the dust into the air. Apply in wall voids, cracks and crevices; in or behind base boards,

service ducts, false floors and ceilings; around electrical entrances, water and sewer pipe entrances; under and behind cabinets, refrigerators, and sinks; in equipment legs and bases; around window and door frames; in attics and crawl spaces.

HOW TO APPLY

- Apply directly into voids, cracks and crevices using hand or power dusters or other suitable equipment. Apply lightly and uniformly.

WHEN TO APPLY

- Every 7 to 10 days, if needed.

RESTRICTIONS

- In living areas, avoid deposits on exposed surfaces and avoid introducing the dust into the air.

- Do not use in aircraft cabins.

- Do not apply in any rooms while the elderly or infirm are present.

- Do not apply in occupied classrooms.

- Do not apply to clothing.

- Do not apply to food/feed areas of commercial food/feed handling establishments, restaurants, or other commercial establishments where food/feed is commercially prepared, processed or served.

- Do not apply to animal feedstuffs or watering equipment.

FOR FOOD/FEED HANDLING ESTABLISHMENTS

TEMPO 1% Dust applications are permitted in non-food/feed areas of food handling establishments only as a void or crack and crevice treatment. It is not permitted in the food/feed areas of food/feed handling establishments. Food/feed handling establishments are defined as places other than private residences where exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also

considered food areas. Non-food/ non-feed handling areas of food/feed handling establishments include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

Limitations, Restrictions, and Exceptions

IN AND AROUND BUILDINGS, STRUCTURES, ETC. - FLYING PEST

APPLICATION FOR PESTS (FLYING PEST)

- Thoroughly dust nests and entrances and surrounding areas. To treat nests in wall voids, drill a hole in the area, dust entrance and surrounding areas. Wear protective clothing if necessary to avoid stings. You may also help avoid stings by treating wasp and bee nests in the evening when the insects are less active and are inside the nest. For best results, check nests within one or two days to ensure complete kill. Remove and destroy treated nests to prevent emergence of newly hatched insects, secondary pest problems, and possible staining from honey. If removal is not possible, retreat the nest, if necessary.

Method

[Directed](#)

[Dusting](#)

Rates

[field rates 0](#)

•

Timings

[When pests are seen or found.](#)