

COMMERCIAL INDOOR TREATMENTS - BOXELDER BUGS, ETC. (0.50% SPRAY)

General Information

PRODUCT INFORMATION

PYROFOS CS is a liquid microencapsulated concentrate containing the active ingredient Chlorpyrifos for use in and around (within 6 to 10 feet) the following sites:

Calf Hutches and Calving Pens and Parlors; Cargo Holds of Ships, Empty Chicken Houses, Food and Non-Food Areas of Manufacturing, Industrial Plants and Food Processing Plants; Hog Barns; Milk rooms; Railroad Boxcars.

PYROFOS CS may be used to treat any surface which will not be damaged or marked by water. Heavy applications may leave a noticeable deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

PYROFOS CS Controls: Angoumois Grain Moths, Aphids, Ants(except carpenter, fire, harvester, and pharaoh), Bees, Beetles, Boxelder Bugs, Cadelles, Carpet Beetles, Cigarette Beetles, Clover Mites, Cluster Flies, Confused Flour Beetle, Crickets, Darkling Beetles, Dermestid Beetles, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Elm Leaf Beetles, Firebrats, Flea Beetles, Flies, Gnats, Grain Mites, Indian meal Moths, Leafhoppers, Leaf Miner, Mealworms, Mediterranean Flour Moths, Millipedes, Moths, Northern Fowl Mites, Psocids, Rod Flour Beetle, Saw-toothed Grain Beetles, Scale, Silverfish, Sowbugs, Spiders(except black widow and brown recluse), Stink Bugs, Trogoderma, and Weevils (Granary Weevils and Rice Weevils).

Limitations, Restrictions, and Exceptions

COMMERCIAL INDOOR TREATMENT

Application within food areas is limited to spot and / or crack and crevice treatments. Even when using as a spot treatment, do not exceed 2 sq. ft. per individual spot. Do not re-apply more than once every 7 days.

NOTE: Use the higher rate for clean-outs, for longer residual control or for high insect infestations.

Apply PYROFOS CS as a crack and crevice residual product where insects may be hiding, living and breeding and/or for selected spot treatment.

COMMERCIAL INDOOR TREATMENTS - BOXELDER BUGS, ETC. (0.50% SPRAY)

Inject into cracks and crevices and/or voids where these insects may be hiding or hibernating. Infested areas include, but are not limited to, cracks and crevices in weep holes, wall voids, around window and door frames, and under siding. Repeat application as necessary. Treat wall area around entry points on exterior where insects gather.

Method

[Injection](#)

Rates

[field_rates 0](#)

•

Timings

[N. A.](#)