

ASPARAGUS

General Information

RESISTANCE MANAGEMENT

CHLORPYRIFOS 15G contains a Group 1B insecticide. Insect and mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by the product or other Group 1B insecticides.

The following practices are recommended to delay development of insecticide resistance:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix product containing insecticides with different modes of action (different insecticide groups), provided the products are registered for use.
- Base insecticide use on comprehensive Integrated Pest Management (IPM) programs.
- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your Local Agricultural Extension Specialist, certified crop advisor, or Manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

RATES OF APPLICATION

Do not apply the product by aircraft at a rate greater than 1 pound chlorpyrifos a.i. (6.6 lbs. of the product) per acre.

Limitations, Restrictions, and Exceptions

ASPARAGUS (For Use only in the Midwest and Pacific Northwest states.)

- Apply the product as a post-harvest ground application for suppression of target pests. Apply as a band over the entire crown area when Asparagus beds have been split (i.e. remove most of the soil from above the crowns). Cover area with soil the day of application. NOTE: Control may be reduced in soils with high organic matter content.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply more than 3 lbs. chlorpyrifos a.i. (20 lbs. of the product) per acre between harvests.

Method

[Broadcast/Foliar Ground](#)

[Band](#)

Pre-Harvest Interval

180 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Exception: If the product is injected or incorporated into the soil, the WPS allows workers to enter the treated area if there will be no contact with anything that has been treated under certain circumstances.

Timings

[Post-harvest](#)