### SOIL APPLICATION

#### General Information

Foliar fertilization is intended to supplement standard ground fertility programs and will not by itself provide all nutrients normally required by agricultural crops.

#### APPLICATION PRECAUTIONS:

#### Tank Mixes

Sysstem-MN is compatible with many fertilizers and pesticides but a compatibility check is always recommended. Fill the tank 1/2 to 2/3 with water and begin agitation and add tank-mix partner products (surfactants, pesticides, etc.) to the tank based on manufacturer labeled instructions. Mixing order will depend on the tank-mix partners. If the tank-mix partners are sensitive to high pH, add Sysstem-MN first to lower the pH of the spray solution. If the tank-mix partners are sensitive to low pH solutions, add Sysstem-MN last and spray immediately after mixing. For adjuvants refer the tank-mix partner labels for the proper mixing order and adjuvant selection. Do not mix Sysstem-MN with fixed copper-based fungicides. Do not apply Sysstem-MN in combination with citrus oils at more than 1 pt/ac or 0.5%, whichever is less or in combination with petroleum oils at more than 2 qts/ac or 1.0%, whichever is less.

## **Spray Applications**

Do not apply Sysstem-MN with any type of hand, pump or back-pack sprayer. Avoid applying to wet leaf and fruit surfaces. Apply to dry plant surfaces to the point of wet with no run-off. Avoid spraying at temperatures above 85 degrees at time of application during day light hours. For additional information regarding Agro-K products, tank-mixes and spray conditions contact your local Agro-K representative. User assumes full responsibility to ensure product, equipment and plant compatibilities.

Limitations, Restrictions, and Exceptions

**Directions for Commercial Use:** 

Soil: Product can be applied at 7 to 14 day intervals for a maximum of 5 times

during the growing season. Can be band sprayed or metered into irrigation system. Product can also be applied as a soil drench on trees.

Method

Soil application

Rates

field\_rates 0

•

# Timings

At 7 to 14 day intervals for a maximum of 5 times during the growing season.