

FOR PERIMETER TREATMENTS

General Information

General Information

This product is a ready-to-use bait for the control of ants, cockroaches, crickets and silverfish including:

- Ants: Argentine, Carpenter, Crazy, Little Black, Odorous House, Pavement, Pharaoh and Thief Ants.
- Cockroaches: American, Asian, Brown, Brown Banded, German, Oriental and Smokey Brown Cockroaches.
- Crickets: Camel, Field, House and Mole Crickets.
- DO NOT contaminate this product with other pesticides. If this product becomes contaminated with another pesticide it will lose its effectiveness.
- DO NOT apply directly to food. In commercial food handling facilities, cover or remove any food during application. In food handling areas, bait may be placed in removable trays. Always record the number and location of bait stations. This product may be used in an Integrated Pest Management program incorporating good sanitation practices.
- This product may be applied in appropriate areas with a power duster (except in California). Consult equipment manufacturer for recommended settings.
- Children and pets should not be in the treatment area until after application is completed.
- DO NOT treat pets with this product.
- Avoid contamination of feed and foodstuffs.
- Avoid contamination of ornamental plants.

Limitations, Restrictions, and Exceptions

FOR PERIMETER TREATMENTS

Spread bait at a rate of 6 ounces or 1.25 cups per 100 square feet in a band at least 2 feet wide around foundations, patios, driveways, sidewalks, entrances to homes and buildings, or other areas where insects are present. Apply around landscape lighting, floodlights, street lamps or other exterior fixtures and scatter bait in flower

beds, leaf litter, wood piles, trash cans and refuse areas. Reapply in seven days if necessary.

As a maintenance treatment apply quarterly applications at a rate of 1 pound per 1,000 square feet around the perimeter and to harborage areas to prevent insect populations from developing around structures in warm months. This product works well in combination with registered nonrepellent insecticides including dinotefuran, chlorfenapyr, and fipronil. If the perimeter is treated with a repellent insecticide, the bait may still be used around the perimeter, but it should be applied in a band outside the repellent barrier for best results.

Method

[Broadcast bait](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Timings

[N.A.](#)