

# **INDOOR USE - PERIDOMESTIC COCKROACHES**

## General Information

## Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## Use Restrictions

- Apply this product only as specified on this label.
- DO NOT apply this product directly to domestic (companion animals) such as dogs, puppies, cats or kittens.
- Except when applying to livestock, DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to livestock, only protected applicators may be in treated area during application.
- Except when applying to livestock, DO NOT enter or allow others to enter treated area until sprays have dried.
- DO NOT apply more than 1 time per day.
- Remove pets, birds and cover fish aquariums before spraying.
- Remove or cover exposed food and drinking water, in the treatment area, before application.
- Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces in the treatment area, or wash them before use.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies or aquatic habitat can occur. DO NOT allow the product to enter any drain during or after application.
- In hospitals and nursing homes, DO NOT use in patient rooms while occupied.
- In schools, DO NOT apply to classrooms when in use.
- DO NOT use in aircraft cabins.
- DO NOT make outdoor applications during rain.
- DO NOT apply directly to impervious horizontal surfaces such as sidewalks, driveways and patios except as a spot or Crack & CreviceR treatment. During application, DO NOT allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters.

- All outdoor applications, if permitted elsewhere on this label, must be limited to spot or Crack & Crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:

1. Applications to soil or vegetation, as listed on this label, around structures;
  2. Applications to lawns, turf and other vegetation, as listed on this label;
  3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
  4. Applications to underside of eaves, soffits, doors or windows permanently protected from rainfall by a covering, overhang, awning or other structure;
  5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed 1 inch in width;
  6. Applications made through the use of a coarse, lowpressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutter or surface waters, in order to control occasional invaders or aggregating pests.
- DO NOT water the treated area to the point of runoff.
  - DO NOT wet plants to point of runoff or drip.
  - DO NOT wet articles to point of runoff or drip.
  - DO NOT use treated articles until spray has dried.
  - When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
  - When used in grain elevators: Apply to floor and up to 15 feet up interior walls of bins and storage with a light, coarse, fan spray. Apply to bin exterior up to 3 feet from ground. Treat a 3 foot border on ground surrounding the bin. Use sufficient water to cover the treated area, DO NOT allow runoff to occur. DO NOT apply directly to grain.

#### Application Instructions

Apply via Crack & Crevice, direct, spot or surface application method(s). DO NOT apply as a space spray.

#### Hose-end Sprayers

Dilute, as desired, according to the dilution table or specific directions for use. Shake sprayer to disperse the product and shake occasionally during use to maintain dispersion.

## Limitations, Restrictions, and Exceptions

### Indoor Use

#### Peridomestic Cockroaches

Apply at the 0.10% rate, as a Crack & Crevice or spot treatment to hiding places such as cracks and crevices, moist areas, openings around pipes and sinks, under refrigerators and manholes. Treat floors, walls, under pallets, behind shelves and drawers. Apply directly on insects in these locations where possible.

#### Method

[Spot treatment](#)

#### Rates

[field\\_rates 0](#)

- 

#### Timings

[N.A.](#)