

NON-CROP SITES - WEEDS CONTROLLED AT 1/2 TO 1 OUNCE PER ACRE

General Information

Do not use on food or feed crops except as specified by this label.

IMPORTANT RESTRICTIONS

DO NOT USE ON FOOD OR FEED CROPS EXCEPT AS SPECIFIED BY THIS LABEL. Injury to or loss of desirable trees or other plants may result if the restrictions listed below are not followed.

- Do not apply Omni Brand MSM 60 DF Herbicide (except as specified), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the product may be washed or moved into contact with their roots.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water.
- Keep from contact with fertilizers, insecticides, fungicides and seeds.

Spraying and mixing equipment used with Omni Brand MSM 60 DF Herbicide must not be used for subsequent applications to food or feed crops with the exception of pastures, rangeland, and wheat, as low rates of Omni Brand MSM 60 DF Herbicide can kill or severely injure most food or feed crops.

Use Omni Brand MSM 60 DF Herbicide only in accordance with directions on this label. Helena Chemical Company will not be responsible for losses or damages resulting from the use of this product in any manner not specifically specified by Helena Chemical Company. User assumes all risks associated with such non-specified use.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.

- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.
- Do not apply this product through any type of irrigation system.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc. Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when the crop canopy is dense. Avoid swath overlapping, and shutoff spray booms while starting, turning, slowing, or stopping to avoid crop injury.

Do not make applications using equipment and/or spray volumes or under weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift, refer to the Spray Drift Management section of the label. Continuous agitation is required to keep Omni Brand MSM 60 DF in suspension.

At the End of the Day

When multiple loads of Omni Brand MSM 60 DF Herbicide are applied, it is recommended that at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of Omni Brand MSM 60 DF Herbicide as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal. of household ammonia* (contains 3% active) for every 100 gal. of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing

cleaning agent and water.

4. Repeat Step 2.

5. Rinse the tank, boom, and hoses with clean water.

6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) specified on this label. Do not exceed the maximum-labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

*Equivalent amounts of alternate-strength ammonia solution or a Helena Chemical Company approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or Helena Chemical Company representative for a listing of approved cleaners.

Notes:

1. Attention: Do not use chlorine bleach with ammonia as dangerous gasses will form. Do not clean equipment in an enclosed area.

2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.

3. When Omni Brand MSM 60 DF Herbicide is tank mixed with other pesticides, examine all required cleanout procedures and follow the most rigorous procedure.

4. In addition to this cleanout procedure, follow all precleanout guidelines on subsequently applied products as per the individual labels.

5. Where routine spraying practices include shared equipment frequently being switched between applications of Omni Brand MSM 60 DF Herbicide and applications of other pesticides to Omni Brand MSM 60 DF Herbicide sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to Omni Brand MSM 60 DF Herbicide to further reduce the chance of crop injury.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of

action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistance weed biotypes, it may be necessary to change cultural practices within and between crop seasons, such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or Herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as a part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

Limitations, Restrictions, and Exceptions

NON-CROP SITES

Application Information

Omni Brand MSM 60 DF Herbicide is for general weed control on private, public and military lands as follows: Uncultivated areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - non-crop producing (such as farmyards, fuel storage areas, fence rows, soil bank land, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.). It is also for the control of certain noxious and troublesome weeds.

Consult the "Weeds Controlled" and "Brush Species Controlled" tables to determine the appropriate application rate.

Omni Brand MSM 60 DF Herbicide may be applied in tank mixture with other Herbicides labeled for use on non-crop sites. Fully read the labels and follow all the

directions and restrictions on each label.

Applications may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the target vegetation with the application equipment being used.

Application Timing

For best results, Omni Brand MSM 60 DF Herbicide should be applied postemergence to young, actively growing weeds. Applications may be made at any time of the year, except when the ground is frozen.

GRASS REPLANT INTERVALS

Following an application of Omni Brand MSM 60 DF Herbicide to non-crop areas, the treated sites may be replanted with various species of grasses at the intervals specified in the label.

The specified intervals are for applications made in the spring to early summer. Because Omni Brand MSM 60 DF Herbicide degradation is slowed by cold or frozen soils, applications made in the late summer or fall must consider the intervals as beginning in the spring following treatment.

Testing has indicated that there is considerable variation in response among the species of grasses when seeded into areas treated with Omni Brand MSM 60 DF Herbicide. If species other than those listed above are to be planted into areas treated with Omni Brand MSM 60 DF Herbicide, perform a field bioassay, or previous experience may be used, to determine the feasibility of replanting treated sites.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)