

VEGETABLES - PEPPERS - BROADCAST SPRAY

General Information

PRODUCT INFORMATION

PRODUCT INFORMATION Triad contains a blend of plant growth regulators acting as a biostimulant. This product is designed to enhance plant growth and development. Triad stimulates nutrient uptake, nutrient utilization, cell division, cell differentiation, and cell enlargement. This product is compatible with pesticides and is particularly effective when applied with fertilizers.

MIXING INSTRUCTIONS: This product readily dissolves in water to form a spray to be applied through conventional liquid application and sprinkler irrigation systems. Shake this product before use. To achieve thorough spray coverage, dilute with a sufficient volume of water, however avoid excessive runoff. Apply within 12 hours of spray solution preparation and be sure to maintain agitation during application. This product may be applied as a tank mix with foliar fertilizers, fungicides, herbicides, and insecticides but should be added last when preparing the spray mixture.

APPLICATION INSTRUCTIONS Apply this product by ground or air to foliage at a finished spray volume of 2-10 gallons per acre, unless specified otherwise on the label. The larger spray volumes of water may be used if it does not create excessive runoff. For best results, make applications in early morning or late evening. If applying this product as a band or foliar-directed spray, the application rate should be decreased from the recommended broadcast rate proportional to the percent of the field surface covered by the spray but DO NOT apply below the minimum rate.

Limitations, Restrictions, and Exceptions

VEGETABLES

PEPPERS

Number of Applications: 4-6

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Exception: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Apply at transplant, or 3-4 leaf stage for direct seeded spray, and then 7-14 day intervals.](#)