

# **APPLES, PEARS, CHERRIES, PLUMS, CITRUS, ALMONDS, NUTS, PEACHES, AVOCADO, FIGS**

## General Information

### DIRECTIONS FOR USE

For the correction of copper deficiencies in crops.

Coating on Fertilizer Blends - This product may be used as a coating on all fertilizer blends (including urea or urea blends). Application rates and timings should be based on a soil and/or tissue test, as well as consultations with a professional agricultural consultant. Apply up to 1 lb of Wolf Trax Nutrient per 100 lbs of fertilizer (1 kg per 100 kg of fertilizer). Combined amounts of all Wolf Trax Nutrients added beyond 1 lb per 100 lbs of fertilizer may not adhere to the fertilizer. To apply, pour the required rate on to the fertilizer as it is being blended. Allow the product to mix thoroughly with the blend for several minutes.

Mixing instructions for foliar applications - Fill tank to 1/3 full, start agitation and pour package contents into tank. Use sufficient water to ensure thorough coverage. If mixing with a crop protection product, always add the micronutrient to the spray solution before adding the crop protection product. Prior to spraying, conduct a compatibility test by adding the products to water in a quart jar to ensure that the products will remain in solution. It is also recommended to use 50 mesh or coarser screens on all nozzles.

Foliar application rates and timing -The rates are suggestions based on correcting deficiencies seen in extensive field-testing. Individual rate recommendations may be obtained using a tissue test.

### Limitations, Restrictions, and Exceptions

#### Foliar Timing

Apply to foliage as a cover spray in spring. Repeat application if necessary up until bloom. (May also be applied to soil in liquid fertilizer at rates of up to 2.5 lbs/acre (2.8 kg/ha))

#### Method

Foliar spray

Rates

field rates 0

field rates 1

- 

Timings

Apply in spring and repeat application if necessary up until bloom.