

REDUCED RATE IN SPRING WHEAT, INCLUDING DURUM WHEAT - MONTANA, NORTH DAKOTA, & SOUTH DAKOTA

General Information

PRODUCT INFORMATION

EVEREST 2.0 HERBICIDE is for use in spring, durum and winter wheat. EVEREST 2.0 HERBICIDE controls wild oat, green foxtail, yellow foxtail, Italian ryegrass, windgrass, barnyardgrass, brome species and numerous broadleaf weeds, including redroot pigweed, wild mustard and shepherd's purse. EVEREST 2.0 HERBICIDE also suppresses additional grass and broadleaf weeds, including downy brome, and wild buckwheat.

EVEREST 2.0 HERBICIDE is absorbed by foliage and roots of susceptible weeds, which cease growth soon after application. Weed emergence is not necessary for control due to the soil residual activity provided by EVEREST 2.0 HERBICIDE. Maximum weed control is achieved one to two weeks after application, though susceptible weeds will stop growing and will no longer be competitive soon after application. For broader spectrum activity, EVEREST 2.0 HERBICIDE may be tank-mixed with a broadleaf herbicide listed on this label. See TANK-MIXES section for recommended products.

RESISTANCE MANAGEMENT

EVEREST 2.0 HERBICIDE is an acetolactate synthase (ALS) inhibitor, and will therefore control weed biotypes which have developed target site resistance to certain classes of herbicides, including ACCase inhibitors, dinitroanilines and triallates.

Any weed population may contain or develop plants naturally resistant to an herbicidal mode of action. Weed populations resistant to ALS inhibiting herbicides already exist. EVEREST 2.0 HERBICIDE will not control ALS resistant weeds. Resistant biotypes may eventually dominate the weed population if herbicides with an identical mode of action are used repeatedly in the same field and weed control may fail. Where possible, rotate the use of EVEREST 2.0 HERBICIDE with herbicides that have a different mode of action.

Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. The use of EVEREST 2.0 HERBICIDE should conform to resistance management strategies established for the use area. Consult your agricultural advisor for resistance management strategies and recommended pest management practices for your area.

Read the entire DIRECTIONS FOR USE before using EVEREST 2.0 HERBICIDE.

This product is not recommended for use on flood irrigated fields.

USE RESTRICTIONS

- For use only in wheat.
- Make only one application per year.
- Do not graze livestock or harvest forage for hay from treated areas for a minimum of 30 days following application.
- Do not mix, load or clean spray equipment within 33 feet of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Do not apply within 50 feet of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Do not apply post emergence when rain is expected within the next hour after application.
- Do not allow this chemical to drift onto other crops.

- Do not harvest grain for 60 days following application.
- Do not apply this product through any type of irrigation system.
- For Idaho, use only in the counties of Benewah, Boundary, Bonner, Clearwater, Idaho, Kootenai, Latah, Lewis, Nez Perce, and Shoshone. Use in all other counties of Idaho is prohibited.

POSTEMERGENCE USE DIRECTIONS

FOR SPRING, DURUM AND WINTER WHEAT

APPLICATION PROCEDURES

GROUND APPLICATION

Apply in a spray volume of 5 to 10 gal/A (or 50 to 100 L/ha) at 30 to 50 psi to ensure proper weed coverage. Use nozzles that provide a medium to coarse size droplet for best coverage and drift control.

AERIAL APPLICATION

Apply in water using a minimum spray volume of 3 gal/A (or 30 L/ha). For best results, use a minimum of 5 gal/A (or 50 L/ha) under dry conditions or heavy weed infestations. Use nozzles that provide 200 to 350 micron size droplets for best results and to insure uniform spray coverage. Aerial applications with EVEREST 2.0 HERBICIDE must be made with low drift nozzles at a maximum height of 10 feet above the crop and at a maximum pressure of 40 psi. Do not apply aerially when wind speed is greater than 10 mph. Do not allow spray to drift onto adjacent crops, as injury or loss may occur.

ENDANGERED SPECIES PROTECTION

To avoid adverse effects on endangered dicot plant species, the following measures will be required where endangered plant species occur in the counties listed in the following table:

State: Idaho

County: Idaho, Lewis, Nez Perce

State: Minnesota

County: Brown, Cottonwood, Goodhue, Jackson, Renville

State:Montana

County:Flathead, Lake

State:Oregon

County:Benton, Clackamas, Lane, Linn, Marion, Polk, Union, Wallowa, Washington, Yamhill

State:Washington

County:Asotin, Chelan, Cowlitz, Lewis, Lincoln, Spokane, Whitman

State:Wyoming

County:Laramie

For ground applications, the applicator must:

- Apply when there is sustained wind away from native plant communities,
- OR
- Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets,
- OR
- Leave a 50-foot untreated buffer between the treatment and native plant communities

For aerial applications, the applicator must:

- Apply only when there is sustained wind away from native plant communities,
- OR
- Leave a 350-foot untreated buffer between the treatment and native plant communities

Limitations, Restrictions, and Exceptions

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of EVEREST 2.0 Herbicide (EPA Reg. No. 66330-391).

EVEREST 2.0 Herbicide can be used at a reduced rate in spring wheat, including durum wheat, for high pH and light texture soils. 0.75 fl oz/A can be used when soil pH is 7.8 or greater and if organic matter is less than 3% for control of green foxtail and wild oat as well as suppression of brome species.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Crop\)](#)