

BANANA, PLANTAIN

General Information

Chemigation: Do not apply this product through any type of irrigation system.

Spray Volume: ORIUS 45DF may be applied in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under "REMARKS" for each crop. Increase spray volume as crop growth increases to ensure thorough coverage of the foliage and fruit.

Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

No aerial application is allowed on Long Island, New York.

Mixing: Add recommended amount of ORIUS 45DF into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the ORIUS 45DF should be thoroughly dispersed prior to the addition of other materials.

Compatibility: To determine the compatibility of ORIUS 45DF with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly, and allow to stand at least five (5) minutes. If the combination remains mixed or can be remixed readily, the mixture is considered physically compatible. For further information, contact your local Makhteshim Agan representative.

Limitations, Restrictions, and Exceptions

Apply specified dosage as a foliar spray in a minimum of 20 gallons or more of water per acre to assure thorough coverage of the plants. A spray oil recommended for use on banana or plantain should be utilized at 4.3 pints per acre in tank-mix with ORIUS 45DF. Thoroughly mix ORIUS 45DF in the spray solution prior to the addition of a spray oil. Make the first application before disease becomes established and repeat applications at 14-day intervals if necessary to maintain

disease control.

Do not apply ORIUS 45DF to non-bagged bananas. A maximum of 16 oz of ORIUS 45DF may be applied per acre per harvest cycle. ORIUS 45DF may be applied up to the day of harvest.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Foliar spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours