

CITRUS: FIELD APPLICATIONS - TANGERINE HYBRIDS

General Information

GENERAL DIRECTIONS FOR USE

Use only as directed. Read the label thoroughly and make sure it is understood before making applications. Keep out of reach of children.

5.1 Application Instructions:

- ProGibb 40% water-soluble granule contains gibberellic acid which is an extremely potent plant growth regulator; when applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, results in undesirable effects. Always consult the Valent Agricultural Specialist in your area for the spray regimen best suited to your conditions.

- Do not apply to plants under pest, nutritional, or water stress.

- When a range of rates is indicated, use the concentration and spray volume indicated locally by the Valent

Agricultural Specialist.

- For optimum effectiveness, thorough spray coverage must be achieved; all parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Discard any unused spray material at the end of each day following local, state or federal law.

- For most efficacious results, the water pH is best at 7.0, and always below 8.5.

- ProGibb 40% applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.

- Product persistence: Re-apply ProGibb 40% if significant rain occurs within 2 hours

of application.

- Compatibility: When considering the tank mixing of ProGibb 40% with other products, use the following compatibility jar test before mixing a whole tank.

Start with a clear glass or plastic quart jar. Add water from the same water source that will be used for the larger tank mix. Add the pesticides in correct proportions. Mix thoroughly and let stand for a minimum 15 minutes. Heat, separation, gelling, are all signs of incompatibility.

Before using any mixes that pass the jar tests for compatibility, it is imperative to test it on a designated area as it may result either in phytotoxicity or ineffectiveness. For further information, consult your Valent agricultural specialist.

- DO NOT apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).

- No preharvest interval is required for this product.

Limitations, Restrictions, and Exceptions

SPRAY GUIDELINES FOR CITRUS

For citrus, apply in sprays of sufficient water volumes to ensure thorough fruit wetting. In most cases, this application will cause some drop of older mature leaves; this drop of older leaves is inconsequential. However, application to trees of low vigor or under stress (pest, nutritional, or water, etc.) causes severe leaf and/or fruit drop. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank. Applications of copper fungicides and/or oils within three weeks (before or after) the ProGibb 40% application often results in significant leaf drop and fruit drop.

To delay disorders associated with rind aging, puffiness, and softening, and to increase peel strength of tangerine hybrids.

Make one spray application two weeks prior to color break. Apply as a dilute spray.

NOTE: Do not apply if early harvest is planned. Do not apply after coloring as pre-harvest rind staining has been known to occur. Application during coloring sometimes causes variation in rind color development.

Method

[Dilute Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[prior to color break](#)