

NONFOOD/ORNAMENTALS - GRASSES (ORNAMENTALS IN NON-PASTURED AREAS ONLY)

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT allow this product to drift.

DO NOT apply this product to seeds or seed pieces.

DO NOT use for aerial application.

DO NOT apply this product with aerial or chemigation equipment.

FOLIAR SPRAY DRIFT MANAGEMENT

Avoiding spray drift from foliar applications is the responsibility of the applicator. The interaction of many equipment-and-weatherrelated factors determine the potential for spray drift from foliar applications. To protect water resources, the applicator and the grower are responsible for considering all these factors when making decisions.

Consult State Agricultural Experiment stations or State Agricultural Extension Service for additional information as the time of applications needed will vary with the local conditions.

COMPATIBILITY AND PLANT SAFETY

CAPTAN 50 WETTABLE POWDER can be combined safely and effectively at listed dosage rates with most commonly used fungicides and insecticides, with the

exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and Bordeaux mixture will reduce the fungicidal activity of captan. Do not apply captan in combination with or immediately before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with captan. The time factor governing the safe interval between captan and oil sprays varies due to general climatic conditions; therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders, which cause excessive wetting, is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, captan may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use captan and other spray materials at lowest listed rates and avoid drenching trees.

For concentrate spray applications, apply the same amount of CAPTAN 50 WETTABLE POWDER per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage.

USE PRECAUTIONS

Except as specified begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favors disease(s). Apply the low rate and/or spray at larger intervals when climatic conditions least favors disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the eastern or western U.S. The dividing line between eastern and western U.S. is the Rocky Mountains unless specifically noted otherwise. Follow those directions for your growing area where specified.

Limitations, Restrictions, and Exceptions

NONFOOD/ORNAMENTALS

USE PRECAUTIONS - Do not apply spray to ornamental plants beyond the point of drip from the leaf surface. Apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot of area.

GRASSES - ORNAMENTALS IN NON PASTURED AREAS ONLY

Do not apply to turfgrass in recreational sites, including homes, apartment buildings, daycare centers, schools, playgrounds, parks, recreational areas, sports fields, or other residential areas. The maximum application rate for golf course turf and turf in commercial settings is 4.3 lb ai/A with a maximum seasonal application rate of 8.6 lb ai/A.

Leafspot, Damping-off, Brown Patch, Melting Out, Seedling Blights, Brown Spot on St. Augustine Grass - Use 2 pounds CAPTAN 50 WETTABLE POWDER per 100 gal of water. Apply 10 gal spray per 1000 sq ft. Begin when growth starts in spring. Do not make more than two applications per season. Do not graze treated areas or feed clippings to livestock.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Restricted Entry Interval

48 hours

Timings

[When growth starts in spring.](#)