

# **CHRISTMAS TREES PLANTATION AND NURSERY - PALES WEEVIL**

## General Information

### GENERAL INFORMATION

Lorsban 50W insecticide in water soluble packets is designed for use as a cover or foliar spray to control various pests. The pests controlled are listed in the accompanying table. Generally, Lorsban 50W is compatible with insecticides, miticides, and fungicides commonly recommended for agricultural crops except for alkaline materials such as Bordeaux mixture and lime. Prepare a small amount of the spray mixture and check for compatibility before mixing a large volume of spray.

### General Use Precautions and Restrictions

- Insect control may be reduced at low spray volumes under high temperature and wind conditions.
- Some reduction in insect control may occur under unusually cool conditions.
- Flood Irrigation: To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours following application of Lorsban 50W.
- Chemigation: Do not apply this product through any type of irrigation system.

Handling Precautions for Water Soluble Packets: Do not remove water soluble packet from the overpack except for immediate use. Do not allow packets to come into contact with water prior to use. Do not handle water soluble packets with wet hands or wet gloves. Do not open water soluble packets. Partial use of packets is not allowed. Carefully reseal package containing unopened water soluble packets and protect package from moisture.

### Application Guidelines

#### Broadcast Foliar Application

Apply with conventional power-operated spray equipment using nozzles and spray pressures recommended for insecticides. Apply Lorsban 50W in a spray volume of not less than 2 gallons per acre (gpa) for aerial application equipment (fixed wing or

helicopter) or not less than 10 gpa for ground equipment, unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy.

Ground Application: Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's recommendations for insecticide nozzles with respect to nozzle type, pressure, and spacing.

#### Broadcast Soil Application

Apply with conventional power-operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets (235-400 microns). Unless otherwise indicated, a spray volume of 10 gpa or more is recommended. For band application, use proportionally less spray volume.

#### Aerial Application

Use a minimum spray volume of 2 gpa and follow recommendations for best management practices for aerial application, above.

Marking of swaths by flagging, permanent markers or use of GPS equipment is recommended.

#### Limitations, Restrictions, and Exceptions

#### CHRISTMAS TREES PLANTATION AND NURSERY

##### Pest-Specific Use Directions:

- For pales weevil control, apply rate as a cut stump drench.

##### Specific Use Precautions:

Phytotoxicity: Do not apply under conditions of extreme heat or drought stress. Environmental factors and varietal differences significantly influence potential phytotoxic expression. Testing has shown that Lorsban 50W may be used at recommended rates on the following conifer species without serious phytotoxicity: balsam fir, concolor fir, Douglas fir, eastern white pine, Fraser fir, grand fir, noble fir, Scotch pine, white spruce. Before treating large numbers of other conifer species, it

is recommended that a small block of plants be treated and observed 7 to 10 days for symptoms of phytotoxicity.

Note: The user assumes responsibility for determining if it is safe to treat other conifer species with Lorsban 50W under commercial growing conditions.

#### Specific Use Restrictions:

- Do not make more than 3 applications of Lorsban 50W or other product containing chlorpyrifos per season.
- Do not make a second application of Lorsban 50W or other product containing chlorpyrifos within 7 days of the first application.
- Do not allow livestock to graze in treated areas.

#### Method

[Cut Stump](#)

[Drench](#)

#### Rates

[field\\_rates 0](#)

•

#### Restricted Entry Interval

24 hours

#### Timings

[N.A.](#)