

CITRUS - TEXAS

General Information

USE PRECAUTIONS

Not for use in residential areas.

Read all precautions and directions before using. Apply this product only as specified on the label.

Imidan 70-W is compatible with most commonly used insecticides and fungicides, but is incompatible with alkaline materials such as spray lime, lime sulfur, and Bordeaux mixtures. These materials will reduce the insecticidal activity of Imidan 70-W.

Insecticidal activity may also be reduced when the spray solution has a pH of 7 or higher. The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum insecticidal activity.

DILUTION DIRECTIONS

The rate required for thorough, uniform coverage varies with plant growth at time of application. Apply recommended rate in adequate spray volumes to provide complete coverage of fruit and foliage.

- Dilute or concentrate applications can be made. Use conventional ground equipment.

- For aerial applications, apply in a minimum of 2 gals. of water per acre for field and row crops, and a minimum of 5 gals. of water per acre for tree and vine crops unless otherwise specified in the recommendation for a specific crop.

RESISTANCE MANAGEMENT

Imidan is an organophosphate insecticide. Based on historical use patterns in some areas, certain pest species listed on the label may have developed resistance to organophosphate insecticides. Consult your local agricultural advisor, State

Cooperative Extension Service, or regional Gowan Company representative for recommendations.

USE LIMITATIONS

Do not exceed the maximum rate of Imidan 70-W per acre or the time limitations specified for the individual crops.

For chemigation application instruction information of Imidan 70-W, please refer to the label.

Refer to the label regarding tank mix information.

Limitations, Restrictions, and Exceptions

CITRUS

RATE: Up to 3 lbs may be used for large trees, high density plantings or applications using high volumes of water.

COMMENTS

Apply as a full coverage spray.

For airblast applications use a minimum of 100 gallons of water per acre.

- Do not use in combination with oils.
- Chemigation: Do not apply this product through any type of irrigation system.
- Please follow the Section 3 label for further application directions.

Method

[Broadcast/Foliar Air](#)

Pre-Harvest Interval

7 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

3 days

Timings

N.A.