

# **BARLEY**

## General Information

### ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

### SEED TREATMENT

This product is a combination of Imidacloprid insecticide and the systemic fungicides Metalaxyl and Tebuconazole. This product aids in the control or suppression of the following insects, seed, seedling and soilborne diseases of Barley and Wheat.

### SEED LABELING

Seed commercially treated with Dyna-Shield Foothold Extra Seed Treatment must be labeled in compliance with all the requirements of the Federal Seed Act.

In addition, all bags containing treated seed must be labeled or tagged with the following information:

The rotational crop restrictions as listed on this label.

"This seed has been treated with Foothold Extra which contains imidacloprid, metalaxyl & tebuconazole."

"Do not use treated seed for feed, food or oil processing."

"Store away from feeds and other foodstuffs."

"Do not allow children, pets or livestock to have access to treated seeds."

"Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed."

"Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

"Dispose of seed packaging in accordance with local requirements."

"Do not graze or feed livestock on treated areas for 45 days after planting."

## Limitations, Restrictions, and Exceptions

Insect Control: Early season protection of seedlings against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid) and Wireworms.

Disease Control: Early season disease control of Pythium damping-off and Covered smut, Loose smut, Barley stripe (suppression), Early season Rhizoctonia root rot, Early season common root rot, Early season Fusarium foot rot, Early season suppression of powdery mildew and Early season suppression of barley leaf rust.

APPLICATION DIRECTIONS: Apply prior to planting as a slurry treatment. This product is to be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time. Ensure thorough coverage. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

RESTRICTIONS: Do not graze or feed livestock on treated areas for 45 days after planting.

Method

[Seed Treatment](#)

[Slurry treatment](#)

Rates

[field rates 0](#)

- 

Restricted Entry Interval

24 hours

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Preplant](#)