

POSTEMERGENCE APPLICATIONS TO ROUNDUP READY CORN 2 - PERENNIAL WEEDS

General Information

GENERAL INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THE LABEL OR CURRENT SUPPLEMENTAL LABELING ISSUED BY MANUFACTURER.

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

Hand-held sprayers may also be used.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of aboveground growth and deterioration of underground plant parts.

Unless otherwise specified on the label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of the label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Reduced or unacceptable control may result if weeds or brush are treated under poor growing conditions such as drought stress, disease or insect damage. Reduced results may also occur when treating weeds or brush heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, it is recommended that a residual herbicide program specified on the label be used. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

Mixing this product with herbicides or other materials not recommended in the label may result in reduced performance. However, unless otherwise prohibited on the label or the label of an intended tank mix product, this product may be applied in combination with any herbicide registered for the same site, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. LIABILITY FOR CROP INJURY, HERBICIDE NONPERFORMANCE OR OTHER LOSS OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THE LABEL, OR SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT, IS SPECIFICALLY DISCLAIMED BY MANUFACTURER. BUYER AND ALL USERS ARE RESPONSIBLE FOR ALL LOSS OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF MIXTURES OF THIS PRODUCT OR OTHER MATERIALS THAT ARE NOT EXPRESSLY RECOMMENDED IN THE LABEL.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs

and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Limitations, Restrictions, and Exceptions

POSTEMERGENCE APPLICATIONS TO ROUNDUP READY CORN 2

GENERAL INFORMATION

Nufarm recommends use of this product only on corn hybrids designated as containing the roundup ready gene.

- Applying this product to corn hybrids which are not designated as Roundup Ready will result in severe crop injury and yield loss.
- The RoundUp Ready designation indicates that the corn contains a patented gene, which provides tolerance to this product.

NOTE: The instructions provided in this supplemental label are specific to, and should only be used with, Roundup Ready Corn 2 hybrids. Do not combine the instructions in this supplemental with any other Roundup Ready corn instructions on labeling for this or other glyphosate containing product.

The use of the higher in-crop over the top rates described in this supplemental label on other than Roundup Ready Corn 2 may cause crop injury and reduced yields.

APPLICATION INSTRUCTIONS

For Roundup Ready Corn 2 from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first, this product may be applied over-the-top broadcast or with drop nozzles. When corn height is 24 to 30 inches (free standing), for optimum spray coverage and weed control, drop nozzles are recommended. For corn heights 30 to 48 inches (free standing) apply this product only using ground application equipment with the drop nozzles adjusted to avoid spraying into the whorls of the corn plants.

Single in-crop applications of this product should not exceed 48 fluid ounces per

acre.

Maximum Allowable Combined Application Quantities Per Season

Preplant, At Planting, Preemergence: Maximum amount of this product which can be applied prior to crop emergence is 5.0 quarts per acre.

Postemergence (in crop): Maximum combined total of multiple in-crop applications from emergence through 48 inch stage is 3 quarts per acre.

Preharvest: Maximum preharvest rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) with 35 percent grain moisture or less until 7 days before harvest is 1 quart per acre. See precautions and restrictions on preharvest applications.

Cropping Season: Combined total per year for all applications may not exceed 8.0 quarts per acre.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought, or when tank mixed with Bullet™ or Micro-Tech™ herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including surfactants, fertilizers and/or micro-nutrients are not recommended with this product since this may result in increased potential for crop injury.

ATTENTION:AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO

DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT. For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 32 fluid ounces per acre. See "WEEDS CONTROLLED" section on the label. AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Weed Control Recommendations

Apply 24 to 32 fluid ounces of this product per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for Nufarm Credit Systemic Herbicide for rate recommendations for specific annual weeds. Nufarm Credit Systemic Herbicide applied at up to 48 fluid ounces per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "WEEDS CONTROLLED" section of the label booklet for Roundup Original herbicide.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting corn.

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Preemergence followed by Postemergence Weed Control Program

USE INSTRUCTIONS: This product may be applied postemergence in-crop following any labeled preemergence herbicide application.

The post application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop.

An in-crop application of this product at the recommended rate will provide control of emerged weeds listed on the label. This product may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready Corn 2 from emergence through the V8 state (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are recommended for optimum spray coverage and weed control. For corn heights 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

Postemergence Only Weed Control Program

USE INSTRUCTIONS: This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready Corn 2 from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are recommended for optimum spray coverage and weed control. For corn height 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

PRECAUTIONS, RESTRICTIONS: Single in-crop applications of this product should not exceed 48 fluid ounces per acre. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage or grain. For applications at preharvest timing (see Preharvest section of this supplemental label), allow a minimum of 7 days between application and harvest or feeding of corn stover or grain. There are no rotational crop restrictions following applications of this product.

Postharvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the

time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)