

## **ROOT VEGETABLES (EXCEPT SUGARBEET) / TUBEROUS AND CORM VEGETABLES - ARMYWORM, COLORADO POTATO BEETLE, ETC.**

### General Information

#### APPLICATION

Dilute DELTA GOLD 1.5 EC in water according to the directions and apply to the target crops at the rates listed. Uniform coverage is essential. Consult your local or state agricultural authorities regarding economic thresholds and timing of application for specific pests. Under heavy pest populations use of the highest labeled rate is recommended.

Do not apply Delta Gold 1.5EC by aerial application in New York State.

DELTA GOLD 1.5 EC may be applied before, during, or after planting. Use the higher labeled rate for soil incorporated applications.

#### RESISTANCE

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

ROOT VEGETABLES (EXCEPT SUGARBEET) / TUBEROUS AND CORM VEGETABLES

## Minimum Gallonage Requirements:

For aerial applications use a minimum of 2 gallons of water per acre. For ground applications use sufficient water to obtain full coverage of foliage.

## NOTES for Use on Root, Tuberous, and Corm Vegetables:

- Allow 3 days between applications of DELTA GOLD 1.5EC.
- Except for radishes, DO NOT apply to root, tuberous, and corm vegetables whose tops or leaves are to be harvested for human or livestock consumption.
- Green peach aphid: Suppression only.
- Armyworm (all species), Colorado potato beetle, Lygus bug (plant bug), Tarnished plant bug: See resistance statement under the "Directions For Use" section.
- Armyworm (all species): For best control apply to 1st and 2nd instar larvae.

## Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

## Pre-Harvest Interval

3 days

## Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

## Restricted Entry Interval

12 hours

## Timings

[N.A.](#)