

SPREADER APPLICATION

General Information

- Combination fertilizer and Dimension® preemergent herbicide.
- Provides preemergent control of crabgrass and other annual grasses and broadleaf weeds.
- Provides early postemergent crabgrass control in limited conditions.
- Consistently sized particles reduce segregation potential
- Small particle formulation more particles per square inch for better herbicide distribution.
- Low nitrogen, high potassium formula.
- High active ingredient concentration to accommodate high herbicide rates.

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfgrass species mentioned in the label that have been determined to be tolerant to applications of this product.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Postemergence crabgrass control applications are only effective on crabgrass if applied before crab-grass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

FLORIDA APPLICATIONS: We recommend that you follow the Florida Green Industries BMP's at http://www.flaes.org/pdf/BMP_Book_final.pdf

FLORIDA APPLICATIONS: We recommend that you follow the Florida Golf Course BMP's at <http://www.flaes.org/pdf/glfbmp07.pdf>

Limitations, Restrictions, and Exceptions

SPREADER APPLICATION:

50 lbs. treats 14,700 sq. ft. (0.33 acre) at the 152 lbs. product/acre - HIGH RATE (3.4 lbs/1,000 sq. ft.)

Method

[Broadcast Application](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)