

# **GOLF COURSES, SOD FARMS AND OTHER AREAS WHERE TURF IS GROWN - PINK SNOWMOLD**

## General Information

### GENERAL INFORMATION

This product is a broad spectrum fungicide with systemic and contact modes of action for preventive and curative applications. For use on golf courses, sod farms and other areas where turf is grown such as athletic fields, cemeteries, parks and commercial turf. Use of this product on home lawns is prohibited

### GENERAL PRECAUTIONS AND RESTRICTIONS

Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry. Use of this product on home lawns is prohibited. Under certain growing conditions, a temporary discoloration of the grass may occur. This causes no harm and will disappear in a short time.

Do not make more than one application at the HIGH rate (290 lbs. product/acre) per season on sod farms and other turf areas except for golf course greens and tees. Repeat applications at the HIGH rate on greens and tees must not be made sooner than 14 days. Repeat applications at the LOW rate (145 lbs. product/acre) per season on all turf sites must not be made sooner than 7 days.

Do not apply more than 1,872 lbs. of product per acre/season on golf course greens; 1,333 lbs. of product per acre/season on golf course tees; and 667 lbs. of product per acre/season on sod farms, golf course fairways and other turf sites per season.

This product may not be applied within 25 feet (ground application) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Sodfarm turf treated before harvest must be mechanically cut, rolled and harvested.

### TURFGRASS DISEASE CONTROL

- Use this product in a preventative and curative disease control program.
- Properly calibrate spreader to ensure thorough coverage.
- Apply after mowing.
- Under conditions optimum for high disease pressure, use the higher rate and the longer interval.
- For optimum turf quality and disease control, use this product in conjunction with turf management practices that promote good plant health.
- If treated area is subject to flooding or unusually heavy rainfall, or if disease is severe or reappears, the area should be treated again.

#### Limitations, Restrictions, and Exceptions

#### TURFGRASS

Apply one application in the late fall before snow cover. Do not apply on top of snow. If snow melt occurs, a repeat application is recommended four to eight weeks or later after first application, depending on timing of snow melt. Make only one application at the High rate on sod farms and other turf areas except for golf course greens and tees.

DISEASE: Pink Snowmold (with snowcover)

#### Method

[Broadcast](#)

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

#### Restricted Entry Interval

24 hours

#### Timings

[In the late fall before snow cover.](#)