

WHEAT AND BARLEY - WILD ONION OR GARLIC

General Information

GENERAL INFORMATION

PHENOXY 088 is a special formulation containing 2,4-D acid. It is more effective than 2,4-D amines for controlling hard-to-kill weeds such as field bindweed, Russian knapweed, Canada thistle, leafy spurge, cattails, tules and nutgrass. PHENOXY 088 should be applied when daytime temperatures are lower than 80 to 85 degrees. Spraying when temperatures are higher may cause rapid foliar burn and decrease the product's effectiveness. Treatment at higher temperatures also increases the possibility of crop injury. **INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.** Crop varieties vary in response to 2,4-D and some are easily injured. Apply PHENOXY 088 only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company or state Agricultural Extension Service for advice.

GENERAL RESTRICTIONS: Do not apply product through any type of irrigation system. Do not use in or near a greenhouse.

PHENOXY 088 contains 0.35 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.75 pounds of a.e. per acre per crop cycle.

Limitations, Restrictions, and Exceptions

Apply 1½ pints when grain is fully tillered and wild garlic or onion plants are small. Apply 2 pints after harvest in the crop stubble. For control of new fall growth of wild onion or garlic, refer to FALLOWLAND use directions.

Restrictions and limitations for use on cereal grains:

- For aerial application on grain, apply in 2 or more gallons of water per acre. For ground application, a minimum of 10-15 gallons of water per acre is recommended for proper spray coverage.
- Do not mix with oil for crop uses.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Do not feed treated straw to livestock if a preharvest treatment or emergency treatment as described above is applied.

CEREAL GRAIN RESTRICTIONS:

- Pre-Harvest interval is 4 days.
- PHENOXY 088 contains 0.35 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.75 pounds of a.e. per acre per crop cycle.

Method

[Spray](#)

Pre-Harvest Interval

4 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

48 hours

Timings

[Apply 1½ pints when grain is fully tillered and wild garlic or onion plants are small.](#)