

WARM SEASON GRASSES - CRABGRASS, FOXTAIL, ETC.

General Information

FEATURES & USE TIPS:

- Combination fertilizer and Pendimethalin preemergent herbicide.
- Provides unsurpassed preemergent control of:

Annual Grassy Weeds:

Crabgrass

Foxtail

Fall panicum

Goosegrass

Barnyardgrass

Poa annua

Dicot Weeds (before they germinate):

Chickweed

Knotweed

Hop clover

Oxalis

Purshlane

Henbit

Prostrate Spurge

Cudweed

Eveningprimrose

- Consistently sized particles reduce segregation potential
- Small particle formulation provides more particles per square inch for better herbicide distribution.
- Nitrogen content includes 40% Poly-S, specially formulated for superior slow release nitrogen feeding.

Seeding or Sprigging intervals after treatment

Delay reseeding or winter overseeding for at least 3 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

Treatment interval after Seeding or Sprigging

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Refer to label for Spreader settings, application and instructions.

Limitations, Restrictions, and Exceptions

Apply a repeat application of 145-175 lbs/acre after 6-8 weeks for extended control or where heavy weed infestations are expected.

Note: For Centipedegrass, 401 lbs/acre rate or less. For Tall Fescue, not for use on greens and tees.

Precautions

Do not use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have

been made if the turfgrass is under severe stress conditions (disease, drought, low temperature- below 40 degrees Fahrenheit, scalping, and excessive soil moisture).

Do not apply to dormant bentgrass and *Poa annua* (turf that is not actively growing or has experienced frost).

When any preemergence herbicide is applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be extensive in low-cut bermudagrass and on sandy soils.

As with most other preemergence herbicides, application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding. Do not apply to bentgrass or *Poa annua* after late summer.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 mph) due to its staining potential. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Method

[Broadcast](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[Prior to weed germination.](#)