

## **FALL APPLICATIONS: COASTAL SOUTH - PROGRAM 3**

### General Information

#### FEATURES & USE TIPS:

- Patented dispersing granule formula
- Contains Dimension turf herbicide
- 90% of the nitrogen is from MUtech-L

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

#### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby

established ornamentals when used according to label directions.

## USE PRECAUTIONS

**Turfgrass Safety:** This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following is are tolerant to this product:

### Cool-Season Grasses

Bentgrass, creeping\* (*Agrostis palustris*)

Bluegrass, Kentucky (*Poa pratensis*)

Fescue, fine\*\* (*Festuca rubra*)

Fescue, tall (*Festuca arundinacea*)

Ryegrass, perennial (*Lolium perenne*)

### Warm-Season Grasses

Bahiagrass (*Paspalum notatum*)

Bermudagrass (*Cynodon dactylon*)

Buffalograss\*\*\* (*Buchloe dactyloides*)

Carpetgrass (*Amopus affinis*)

Cenfpedegrass (*Eremocbloa ophiuroides*)

Kikuyugrass (*Pennisetum clandestinum*)

St. Augustinegrass (*Stenotaphrum secundatum*)

Zoysiagrass (*Zoysia japonica*)

- Bentgrass, creeping : Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey1', 'Carmen1', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.

- Fescue, fine :Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgarf', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Kokef', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and 'Waldina'. Not all varieties of fine fescue have been tested.

- Buffalograss : DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike- or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these

cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around fruit trees. Keep people and pets off treated areas until dust has settled.

#### Limitations, Restrictions, and Exceptions

##### Fall Applications:

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the \"Program 3\" rates) or suppress (at the \"Program 2\" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - *Poa annua*, bittercress, chickweed, pineappleweed, hen-bit, geranium (Carolina), parsley-piert, shepherdspurse.

##### Method

[Broadcast](#)

##### Rates

[field\\_rates 0](#)

•

##### Timings

[Preemergence \(Weed\)](#)

[Postemergence \(Weed\)](#)