

CHERRIES - (BLACK CHERRY APHID, FRUIT TREE LEAFROLLER)

General Information

GENERAL INFORMATION

MALATHION 5 may be applied by ground sprayers or by airplanes, mixing the MALATHION 5 in water to form an emulsion spray solution. To make the spray solution, half fill the spray tank with water, agitate, add the proper amount of MALATHION 5 and then fill the spray tank with water while still agitating. Continue to agitate during use.

When applying with ground sprayers, calibrate so as to apply the recommended amount of MALATHION 5 in not less than 3 gallons of water per acre. Airplanes should apply the recommended amount of MALATHION 5 in not less than 1 gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

The dosages given under DIRECTIONS FOR USE are the amounts of MALATHION 5 to be applied per acre treated per application. Repeat applications as necessary, unless otherwise directed in DIRECTIONS FOR USE.

Where a dosage range is given, use the lower rate on small or young plants or with light insect infestations; use the higher rate on large or more mature plants or with heavy insect infestations. The hazards and precautions for handling MALATHION 5 in this container are equally applicable to it after dilution with water for spray application.

Limitations, Restrictions, and Exceptions

CHERRIES - (BLACK CHERRY APHID, FRUIT TREE LEAFROLLER);

Cherries: Not registered in California

The rates for use on cherries are based on a standard of 400 gallons per acre dilute spray for mature trees. Do not apply more than 8 lbs. of actual MALATHION 5 per acre to cherry trees. Injury may occur on certain varieties of sweet cherries, particularly in the Northwest.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[N.A.](#)