

HOUSE MICE

General Information

USE RESTRICTIONS: This product may be used to control Norway rats, roof rats, and house mice only at the Urban and Non-urban sites identified below. Do not use this product in sewers. Do not place bait near to or inside of ventilation duct openings. Do not contaminate water, food, feed, food or feed handling equipment, milk or meat handling equipment, or any area where there is a possibility of contaminating food, feed, milk, or meat, or surfaces that come into contact with food, feed, milk, or meat. All placements of this product in USDA-inspected facilities must be in tamper-resistant bait stations. Do not broadcast bait. Wear water-resistant gloves when handling bait.

Urban Areas: This product may only be used in and around homes, industrial, commercial, and public buildings, and similar man-made structures; in transport vehicles (ships, trains, and aircraft) and in and around related port or terminal buildings; and in alleys. All outdoor bait placements must be made in tamper-resistant bait stations, locked and secured as necessary, unless children, pets, domestic animals, and nontarget wildlife are reliably denied access to the bait placement locations.

Non-Urban Areas: This product may only be used inside of homes, and agricultural buildings.

SELECTION OF TREATMENT AREAS: Determine areas where rats will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where rats or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

Limitations, Restrictions, and Exceptions

House Mice: Apply $\frac{1}{4}$ to $\frac{1}{2}$ ounce of bait per placement, using tamper-resistant bait stations as required under "IMPORTANT". Space placements at intervals of 8 to 12 feet in infested areas pertaining to structures. Larger quantities (up to 2 ounces) may be made in areas of extremely high mouse activity. Maintain an uninterrupted

supply of fresh bait for at least 15 days or until fresh signs of feeding by mice cease to appear.

Follow-up: Replace contaminated or spoiled bait immediately. Using water-resistant gloves, collect and properly dispose of dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

Method

[Bait](#)

Timings

[Until fresh signs of feeding by mice cease to appear.](#)