

## **TABLE GRAPES (SEEDED) - CALMERIA, CHRISTMAS ROSE, EMPEROR, RED GLOBE, ROGUE, QUEENS (SIZING)**

### General Information

#### GENERAL INFORMATION

- Use only as directed. Read thoroughly and understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See Conversion Table at end of Directions for Use section.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Ensure that the pH of the water is less than 8.5.
- Absorption of FALGRO 4L into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Re-apply FALGRO 4L if significant rain occurs within 2 hours of application.
- FALGRO 4L has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

### Limitations, Restrictions, and Exceptions

#### SPRAYING GUIDE FOR GRAPES

Application to grapes is best made by ground sprayer. Spray volume, quality and direction are chosen to ensure thorough wetting of all flower clusters and berries.

## TABLE GRAPES

Application of FALGRO 4L for 'Sizing' produces larger berries and clusters when used in conjunction with girdling and thinning techniques.

Timing: 1 spray 14 days after shatter. Berry diameters of approx. 10-15 mm. May increase berry size.

Target Diameter:

- Calmeria, Emperor, Rogue, Christmas Rose, Red Globe: 12-16mm
- Queens: 12-15mm

Seeded Varieties - Conduct trials with small numbers of plants using recommended rates to determine the optimum rates for the situation before applying to the crop.

Rate - A rate of 40-50 ppm/ai is used when dipping or spraying the cluster directly.

Method

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[1 spray 14 days after shatter.](#)