

CITRUS CANKER CONTROL

General Information

Spectrasol Disinfectant is a phosphate-free, pH neutral formulation designed to provide effective cleaning, deodorizing, and disinfection for hospitals, nursing homes, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, athletic facilities, sports stadiums, amphitheaters and convention centers where housekeeping is of prime importance in controlling cross-contamination from treated surfaces.

Spectrasol Disinfectant, when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: Floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, artificial turf surfaces, vanity tops, shower stalls, bathtubs, and cabinets.

Spectrasol Disinfectant is for use in:

- Food processing plants, dairy farms, hog farms, poultry and turkey farms and egg processing plants.
- Veterinary clinics, animal life science laboratories, kennels and cages, breeding and grooming establishments, pet animal quarters, zoos, pet shops, and other animal care facilities.
- For larger areas such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfecting.

Spectrasol Disinfectant is a one-step neutral disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, HIV-2, HBV and HCV) and inhibits the growth of mold and mildew and their odors when used as directed.

Limitations, Restrictions, and Exceptions

Citrus Canker Control:

For prevention of Citrus Canker Disease through treatment of equipment. Effective against *Xanthomonas axonopodis* pv. *citri* (Citrus Canker) at 2000 ppm active quat.

Treat all trucks, vehicles, and equipment thoroughly at a dilution ratio of 1:54 (2000 ppm active quat). Treatment can be applied to trucks, attached trailers, field harvesting equipment; including cargo area, wheels, tires, under carriage, hood, roof, fenders and any other part of transportation equipment that can be taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using maximum rate 2000 ppm quat. After use, all surfaces which come in contact with food or crop must be rinsed with potable water.

Method

[Dip](#)

[Spray](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)