

# **GRASSES: GOLF COURSE TEES, GREENS, AND ORNAMENTAL TURF - (PINK SNOW MOLD)**

## General Information

### APPLICATION INSTRUCTIONS

Quali-Pro Chlorothalonil 500 ZN, a flowable product containing chlorothalonil, is recommended for use as a spray for the control of many important plant diseases.

### RESISTANCE MANAGEMENT

To avoid the development of tolerant or resistant strains of fungi, Quali-Pro Chlorothalonil 500 ZN should always be tank mixed with a fungicide of different chemistry, and/or a fungicide of different chemistry should be alternated with Quali-Pro Chlorothalonil 500 ZN at each application. If after using Quali-Pro Chlorothalonil 500 ZN as recommended and the treatment is not effective, a tolerant or resistant strain of fungi may be present. Discontinue the use of Quali-Pro Chlorothalonil 500 ZN for at least one season.

Quali-Pro Chlorothalonil 500 ZN is effective for use in programs that attempt to minimize disease resistance to fungicides. Quali-Pro Chlorothalonil 500 ZN has a multi-site mode of action and may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of Quali-Pro Chlorothalonil 500 ZN in programs that seek to minimize the occurrence of disease resistance to other fungicides.

### GENERAL PRECAUTIONS

Quali-Pro Chlorothalonil 500 ZN can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control. Do not combine Quali-Pro Chlorothalonil 500 ZN in a spray tank with pesticides, surfactants, or fertilizers, unless prior use has shown the combination to be physically compatible, effective, and noninjurious under your conditions of use. Do not combine Quali-Pro Chlorothalonil 500 ZN with DiPel 4L, Foil, Triton AG-98, Triton B-1956, or Latron B-1956 as phytotoxicity may result from the combination when applied to crops listed

on the label.

Note: Prior to pouring, slowly invert container several times to assure uniform mixture.

Dosage rates on the label indicate pints of Quali-Pro Chlorothalonil 720 SFT per acre, unless specified otherwise. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

## APPLICATION PRECAUTIONS AND REQUIREMENTS

This product must not be applied within 150 feet for aerial and air-blast applications or 25 feet for ground applications of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

### Limitations, Restrictions, and Exceptions

#### GRASSES: GOLF COURSE TEES, GREENS, AND ORNAMENTAL TURF - (PINK SNOW MOLD)

In areas where pink snow mold (*Gerlachia* or *Fusarium* patch) is likely to occur, apply Quali-Pro Chlorothalonil 500 ZN at the recommended rate given in combination with products containing iprodione at 2 oz. active ingredient per 1000 sq. ft. of turf area.

#### MAXIMUM APPLICATION RATE PER YEAR:

- Ornamental turf: 18.2 fl. oz./1000 sq. ft.
- Tees: 36.5 fl. oz./1000 sq. ft.
- Greens: 51.4 fl. oz./1000 sq. ft.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spray](#)

Rates

[field\\_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Before snow cover in autumn](#)