

# **TREE CROPS: CONIFERS (PINES, SPRUCES) - (BOTRYTIS SEEDLING BLIGHT; PHOMA TWIG BLIGHT)**

## General Information

### APPLICATION INSTRUCTIONS

Quali-Pro Chlorothalonil 720 SFT is a flowable product containing chlorothalonil to be used as a spray for the control of many important plant diseases.

### RESISTANCE MANAGEMENT

To avoid the development of tolerant or resistant strains of fungi, Quali-Pro Chlorothalonil 720 SFT should always be tank mixed with a fungicide of different chemistry, and/or a fungicide of different chemistry should be alternated with Quali-Pro Chlorothalonil 720 SFT at each application. If after using Quali-Pro Chlorothalonil 720 SFT as labeled and the treatment is not effective, a tolerant or resistant strain of fungi may be present. Discontinue the use of Quali-Pro Chlorothalonil 720 SFT for at least one season.

Quali-Pro Chlorothalonil 720 SFT is effective for use in programs that attempt to minimize disease resistance to fungicides. Quali-Pro Chlorothalonil 720 SFT has a multi-site mode of action and may be used to delay or prevent the development of resistance to single-site fungicides.

Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of Quali-Pro Chlorothalonil 720 SFT in programs that seek to minimize the occurrence of disease resistance to other fungicides.

### GENERAL PRECAUTIONS

Quali-Pro Chlorothalonil 720 SFT can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control. Do not combine Quali-Pro Chlorothalonil 720 SFT in a spray tank with pesticides, surfactants, or fertilizers, unless prior use has shown the combination to be physically compatible, effective, and noninjurious under your conditions of use. Do not combine Quali-Pro Chlorothalonil 720 SFT with DiPel 4L, Foil ,Triton AG-98, Triton B-1956 as phytotoxicity may result from the combination when applied to crops listed on the

label.

Note: Prior to pouring, slowly invert container several times to assure uniform mixture.

The required amount of Quali-Pro Chlorothalonil 720 SFT should be added slowly into the spray tank during filling. With concentrate sprays, premix the required amount of Quali-Pro Chlorothalonil 720 SFT in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Dosage rates on the label indicate pints of Quali-Pro Chlorothalonil 720 SFT per acre, unless specified otherwise. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

#### APPLICATION PRECAUTIONS AND REQUIREMENTS FOR AGRICULTURAL USE SITES ONLY

This product must not be applied within 150 feet for aerial and air-blast applications, or 25 feet for ground applications of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

#### Limitations, Restrictions, and Exceptions

TREE CROPS: CONIFERS (PINES, SPRUCES) - (BOTRYTIS SEEDLING BLIGHT; PHOMA TWIG BLIGHT)

Apply Quali-Pro Chlorothalonil 720 SFT in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Quali-Pro Chlorothalonil 720 SFT may be applied with aircraft. When concentrate sprays are used or when treating immature trees, the lower rate of Quali-Pro Chlorothalonil 720 SFT listed may be used. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See the following instructions for application and calibration.

DO NOT allow livestock to graze treated areas.

The following spray volumes are recommended as gallons of spray per acre:

Forest Stands

Dilute: Not Used.

Concentrate: 10-20 gallons / acre (aircraft).

Christmas Trees

Dilute: 100 gallons / acre.

Concentrate: 10-50 gallons / acre (aircraft or ground equipment).

Nursery Beds

Dilute: 100 gallons / acre.

Concentrate: 5-10 gallons / acre (ground equipment only).

**FOR TREE CROPS: CONIFERS (PINES, SPRUCES) - (BOTRYTIS SEEDLING BLIGHT; PHOMA TWIG BLIGHT)**

Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7- to 14-day intervals as long as disease favorable conditions persist.

The minimum retreatment interval for established trees is 21 days. The minimum

retreatment in nursery beds is 7 days.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[when cool, moist conditions favor disease development](#)

[When seedlings are 4 inches tall](#)