

# **LEGUME CROPS - PHYTOPHTHORA AND PYTHIUM SPP., ETC. (PRE-PLANT NURSERY APPLICATION)**

## General Information

### GENERAL APPLICATION INSTRUCTIONS

Apply this product by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation and bare root dip. For foliar sprays, apply this product with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage.

Make applications prior to disease development in conjunction with good cultural management practices. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not make foliar applications to plants treated with copper-based compounds at less than 20 day intervals unless instructed to do so by your crop consultant. Allow foliage to dry completely after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

See label for tank mix information.

### Limitations, Restrictions, and Exceptions

### LEGUME CROPS

#### Disease Prevention:

Low Disease Pressure: Repeat applications at 1-2 week intervals.

High Disease Pressure: Repeat applications at 7-10 day intervals.

Suppression of Fusarium and Rhizoctonia: Not for use in California unless accompanied by a supplemental label.

Pre-plant Nursery application: Apply 1 qt/100 gallons water to nursery plants in seedling trays 1-7 day prior to out planting.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

- 

Restricted Entry Interval

4 hours

Timings

[Preplant](#)