

STRAWBERRY - POWDERY MILDEW (GROUND APPLICATION)

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply this product by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation and bare root dip. For foliar sprays, apply this product with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage.

Make applications prior to disease development in conjunction with good cultural management practices. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not make foliar applications to plants treated with copper-based compounds at less than 20 day intervals unless instructed to do so by your crop consultant. Allow foliage to dry completely after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

See label for tank mix information.

Limitations, Restrictions, and Exceptions

STRAWBERRY

Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1-2 week intervals.

High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals.

Suppression Powdery Mildew: Not for use in California unless accompanied by a supplemental label.

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

4 hours

Timings

[At the first onset of the disease.](#)