

KIWI, OLIVES AND TREE NUT CROPS - ROOT ROTS, CROWN ROTS, ETC. (ROOT DIP)

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply this product by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation and bare root dip. For foliar sprays, apply this product with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage.

Make applications prior to disease development in conjunction with good cultural management practices. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not make foliar applications to plants treated with copper-based compounds at less than 20 day intervals unless instructed to do so by your crop consultant. Allow foliage to dry completely after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

See label for tank mix information.

Limitations, Restrictions, and Exceptions

KIWI, OLIVES AND TREE NUT CROPS

Apply as pre-plant dip to the roots for 30 seconds. Plant within 48 hours after dipping. Mix a fresh solution daily.

Suppression of foliar bacterial and fungal diseases, Anthracnose (*Collectotrichum*), Hull rot (*Monilia* spp.), Flower diseases (*Cladosporium* spp.), Alternaria leaf spots (*Alternaria* spp.), raceme blight (*Phytophthora*) in Macadamia.: Not for use in California unless accompanied by a supplemental label.

Method

[Dip](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

4 hours

Timings

[Preplant](#)