

CONIFERS - SUDDEN OAK DEATH (BARK APPLICATION)

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply this product by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation and bare root dip. For foliar sprays, apply this product with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage.

Make applications prior to disease development in conjunction with good cultural management practices. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not make foliar applications to plants treated with copper-based compounds at less than 20 day intervals unless instructed to do so by your crop consultant. Allow foliage to dry completely after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

See label for tank mix information.

Limitations, Restrictions, and Exceptions

CONIFERS IN COMMERCIAL NURSERIES, LANDSCAPE TREES, PLANTATIONS AND FORESTRY AND PARK APPLICATIONS

Inject (according to injection equipment instructions) 0.5 fl oz (15 ml) of the diluted Fungi-Phite solution into one injection hole.

Dosage: One (1) injection per yard (or meter) of canopy diameter at the drip line. (i.e. a tree with 6 yards (or meters) of canopy diameter will require 6 injections via 6 separate injection holes).

Apply the Fungi-Phite solution from the base of the tree to approximately 5 feet (1.5 M) above ground, ensuring complete bark coverage. Spray over active lesions if they are accessible.

Application Rate: Mix Fungi-Phite and water in a 1:1 ratio (i.e. 1 gallon of Fungi-Phite

plus 1 gallon of water). Add to this 3 oz of a silicon based surfactant.

Method

[Bark Application](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[Curative: Inject 2 times per year in the Spring and Fall.](#)

[Preventative: Inject one time in the Spring and Fall.](#)