

## **AGRICULTURAL CROPS, ORNAMENTAL NURSERIES, BUSINESS, ETC.**

### General Information

### DIRECTIONS FOR USE

It is CRITICAL that ISOMATE-LBAM PLUS is applied as directed.

Application: Dispensers must be applied uniformly throughout the treated areas to obtain a reduction in mating. Twist dispensers loosely around host plant branches, vines, etc., in the top part of or slightly above plant canopy.

Note: Isomate LBAM Plus suppresses mating of target pests. Immigration of mated female moths of this species from adjacent external sources of infestation will reduce the level of control.

### Manage by:

- Treatment of external sources of infestation with Isomate LBAM Plus.
- Treatment of external sources of infestation with an effective insecticide.
- Treatment of pheromone treated orchard with insecticide.

Isomate LBAM Plus is best used in large (> 10 acres), uniform and contiguous areas. Supplementary applications of insecticide are strongly advised when Isomate LBAM Plus is used in areas with high populations. All pests must be monitored so that timely intervention with insecticides is possible.

### Limitations, Restrictions, and Exceptions

### AGRICULTURAL CROPS, ORNAMENTAL NURSERIES, BUSINESS, ETC.

### Rate:

- 200-300 dispensers per acre (1.44-2.15 fl oz or 37.06-55.59 g a.i. per application). Use higher rates when pest pressure is high. Do not exceed 150 g a.i. (or 809 dispensers) per acre per year.

Method

[N.A.](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

- 

Timings

[Prior to light brown apple moth emergence in the spring.](#)