

# **PEACHES, NECTARINES, APRICOTS, PLUMS, CHERRIES AND OTHER STONE FRUITS, ETC.**

## General Information

### DIRECTIONS FOR USE

It is CRITICAL that ISOMATE-OFM TT is applied as directed.

Application - Place dispensers in lateral branches in upper third of tree canopy.

Timing - Apply prior to moth emergence in the spring. Consult your local pest control advisor or Pacific Biocontrol representative for proper timing or estimated dispenser longevity in your area.

Note - Isomate-OFM TT suppresses mating of target pests. Immigration of mated female moths of these species from adjacent external sources of infestation will reduce the level of control.

Manage by:

- Treatment of external sources of infestation with Isomate-OFM TT.
- Treatment of external sources of infestation with an effective insecticide.
- Treatment of pheromone treated orchard with insecticide.

Areawide application is most effective. Supplementary applications of insecticide are advised when Isomate-OFM TT is used in orchards with high pest populations. All pests must be monitored so that timely intervention with insecticides is possible.

### Limitations, Restrictions, and Exceptions

PEACHES, NECTARINES, APRICOTS, PLUMS, CHERRIES AND OTHER STONE FRUITS, ETC.

Rate - Minimum of 100 dispensers per acre (1.84 fl oz or 48.0 g a.i. per application). Do not exceed 150 g a.i. (or 312 dispensers) per acre per year.

### Method

N.A.

Rates

field\_rates 0

field\_rates 1

field\_rates 2

•

Timings

Prior to moth emergence in the spring.