

CONTROL OF HOUSE MICE

Limitations, Restrictions, and Exceptions

RATS AND MICE

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in “Urban Areas” and “Non-urban Areas” that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Do not use this product to control *Peromyscus* spp. mice or within the occupied habitats of threatened or endangered *Peromyscus* spp. mice or other types of threatened or endangered rodents. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

URBAN AREAS: To control Norway rats, roof rats and house mice this product may be used in and around homes and industrial, commercial and public buildings; in transport vehicles (ships, trains and aircraft) and in and around related port or terminal buildings. Applications “around” buildings must be made in tamper-resistant bait stations situated along walls or other external parts of buildings (e.g., doorways, ramps and loading docks) where rats or mice might seek to gain entrance.

NON-URBAN AREAS: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings.

SELECTION OF TREATMENT AREAS: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

House Mice: Open pack and apply $\frac{1}{4}$ to $\frac{1}{2}$ ounce of bait per typical placement. Larger placements – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements of the label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer

are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Method

[Bait](#)

Rates

[field rates 0](#)

-

Timings

[N.A.](#)