

# **TURF AND HYDROSEEDING**

## General Information

### AGRICULTURAL CROPS

#### USE RESTRICTIONS:

Use this product to limit damage caused by birds on ripening crops. This product must be applied using appropriate spray equipment. Do not apply when crop is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage. Do not apply through any type of irrigation system.

#### COMPATIBILITY WITH OTHER PESTICIDES:

Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your representative for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the crop.

#### DILUTION DIRECTIONS:

Shake or stir Rejex-it Migrate container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. For ground application, mix Rejex-it Migrate with water at a ratio of 1 part product to 10-20 parts water. For aerial application, mix Rejex-it Migrate with water at a ratio of 1 part product to 1-3 parts water.

#### APPLICATION DIRECTIONS:

- Apply spray mixture so that crop receives thorough coverage.
- Repeat treatment if significant bird damage begins to occur.
- This product may be applied up to the day of harvest

#### Limitations, Restrictions, and Exceptions

## TURF AND HYDROSEEDING

### USE RESTRICTIONS:

Use this product to repel birds from golf courses, other turf areas (including around agricultural buildings and farmsteads), and hydroseeded areas. This product must be applied using appropriate spray equipment. Do not apply when grass is wet or rain is expected.

### DILUTION DIRECTIONS:

Shake or stir Rejex-it Migrate container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix Rejex-it Migrate with water at a ratio of 1 part product to 20 parts water. For example, mix 2 pints of product with 5 gallons of water to make 5.25 gallons of spray mixture.

### APPLICATION DIRECTIONS:

Apply at a rate of 1.2 gallons of spray mixture (0.07 lbs. a.i.) per 1000 sq. ft. of turf. Spray evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application in 4 days or as warranted by Canada goose activity. Do not mow treated area for several days after application.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[N.A.](#)