

FOR FOLIAR APPLICATION TO GRAPES GROWN IN RESIDENTIAL AREAS

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

SEE INDIVIDUAL SITES FOR SPECIFIC POLLINATOR PROTECTION APPLICATION RESTRICTIONS. IF NONE EXIST UNDER THE SPECIFIC SITE, FOR OUTDOOR FOLIAR APPLICATIONS, FOLLOW THESE APPLICATION DIRECTIONS.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

NON-AGRICULTURAL USES

Do not apply PROKOZ ZENITH 2F Insecticide while bees are foraging. Do not apply PROKOZ ZENITH 2F Insecticide to plants that are flowering. Only apply after all flower petals have fallen off.

PROKOZ ZENITH 2F Insecticide can be used as directed on sod farms, and on turfgrass, and ornamentals on such sites: residential home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields.

PROKOZ ZENITH 2F Insecticide can be used by soil application as directed on trees, shrubs, flowers, and groundcovers in sites: in and around the perimeter of industrial and commercial buildings and residential areas, and municipal, city, state, national, and private wooded and forested areas.

RESTRICTION: Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.

RESISTANCE: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. PROKOZ ZENITH 2F Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

Limitations, Restrictions, and Exceptions

DIRECTIONS FOR FOLIAR APPLICATION TO GRAPES GROWN IN RESIDENTIAL AREAS

Apply specified dosage as a foliar spray using 200 gallons of water per acre.

RESTRICTIONS

- Follow application restrictions for Non-Agricultural Uses on page 3 to protect bees and other insect pollinators.
- Do not apply more than a total of 6.0 fl oz of PROKOZ ZENITH 2F Insecticide per acre per year.
- Allow at least 14 days between applications.
- Applications may be applied up to and including day of harvest.
- Keep children and pets off treated area until dry.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

[field rates 3](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N. A.](#)