

ARTICHOKES - ARTICHOKE PLUME MOTH

General Information

DIRECTIONS FOR USE

METHOD OF APPLICATION: To disrupt mating of the artichoke plume moth on artichokes and other crops where the artichoke plume moth is a pest apply CheckMate APM-F at 3.48 [3.5] fluid ounces (103 ml) product per acre (10 g a.i./acre) to 6.97 [7] fluid ounces (206.31 ml) product per acre (20 g a.i./acre). Do not exceed 52.25 fluid ounces (1545.25 ml) product/acre/year (150 g a.i./acre/year).

Ground Equipment: Apply CheckMate APM-F in sufficient water for uniform coverage. Do not apply to point of runoff. Apply to foliage directly.

Aerial Equipment: Apply CheckMate APM-F in sufficient water for uniform coverage.

APPLICATION NOTES: Reapply as needed based on insect monitoring and field scouting programs. The use of approved stickers improves the adherence to the leaf surface and protection from rainfall. In the event of rain, re-treatment will be required if approved stickers are not used. This product is not to be used in fields with overhead irrigation. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the field must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season can prove necessary to provide adequate protection to the developing crop. These supplemental insecticide applications are to be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

[field_rates 3](#)

-

Restricted Entry Interval

4 hours

Timings

[N.A.](#)