

COTTON - PINK BOLLWORM MOTH

General Information

DIRECTIONS FOR USE

METHOD OF APPLICATION: For use on cotton. To disrupt mating of the pink bollworm moth, apply CheckMate PBW-F at 0.309 fluid ounce (9.14 ml) product per acre (2 g/a.i./acre) to 1.546 fluid ounces (45.72 ml) product per acre (10 g/a.i./acre). Do not exceed 23 fluid ounces (680.2 ml) product/acre/year (150 g/a.i./acre/year). Begin application at pinhead square stage of plant growth.

Ground Equipment: Apply CheckMate PBW-F in sufficient water for uniform coverage. Do not apply to point of runoff. Apply to foliage directly.

Aerial Equipment: Apply CheckMate PBW-F in sufficient water for uniform coverage.

APPLICATION NOTES: This product only affects male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the field must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection to developing bolls. These supplemental insecticide applications should be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae. Reapply as needed, based on insect monitoring and field scouting programs.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

-

Restricted Entry Interval

4 hours

Timings

[Begin application at pinhead square stage of plant growth.](#)