

FOR USE ON TOMATOES AND EGGPLANTS

General Information

DIRECTIONS FOR USE

METHOD OF APPLICATION: To disrupt mating of the tomato pinworm on tomatoes and eggplants, apply CheckMate TPW-F at 1.2 fluid ounces (35 ml) product per acre [8 g a.i./acre] to 6.0 fluid ounces (174.7 ml) product per acre [40 g a.i./acre]. Do not exceed 22 fluid ounces (655 ml) product/acre/year [150 g/a.i./acre/year].

Ground Equipment: Apply CheckMate TPW-F in sufficient water for uniform coverage. Do not apply to point of runoff. Apply to foliage directly.

Aerial Equipment: Apply CheckMate TPW-F in sufficient water for uniform coverage.

APPLICATION NOTES

Reapply as needed based on insect monitoring and field scouting programs. For best results, apply early in the season, after detection of the first tomato pinworm moth with a pheromone trap, such as BioLure. In all cases, make the first application two weeks after transplanting tomato seedlings or within 30 days after direct seeding tomatoes. This product only affects male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the field must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection to developing fruit. These supplemental insecticide applications should be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

-

Timings

[Apply early in the season.](#)