

COMMERCIAL GREENHOUSES AND NURSERIES - BROADCAST

General Information

USE DIRECTIONS

Solera Diquat Landscape & Aquatic Herbicide is a nonvolatile herbicidal chemical for use as a general herbicide to control weeds in commercial greenhouses and nurseries; ornamental seed crops (flowers, bulbs, etc. – except in the state of California); landscape, industrial, recreational, commercial, residential, and public areas; turf renovation (all turf areas except commercial sod farms); dormant established turfgrass (bermudagrass, zoysiagrass – nonfood or feed crop); and aquatic areas. Absorption and herbicidal action is usually quite rapid with effects visible in a few days. Solera Diquat Landscape & Aquatic Herbicide controls weeds by interfering with photosynthesis within green plant tissue. Weed plants should be succulent and actively growing for best results. Rinse all spray equipment thoroughly with water after use. Avoid spray drift to crops, ornamentals, and other desirable plants during application, as injury may result. Application to muddy water may result in reduced control. Minimize creating muddy water during application. Use of dirty or muddy water for Solera Diquat Landscape & Aquatic Herbicide dilution may result in reduced herbicidal activity. Avoid applying under conditions of high wind, water flow, or wave action.

Limitations, Restrictions, and Exceptions

COMMERCIAL GREENHOUSES AND NURSERIES - BROADCAST

For general weed control in commercial greenhouses (beneath benches), field grown and container stock, and other similar areas, Solera Diquat Landscape & Aquatic Herbicide may be applied preplant or postplant preemergence in field grown ornamental nursery plantings or postemergence as a directed spray. Solera Diquat Landscape & Aquatic Herbicide may also be applied preemergence in ornamental seed crops (except in the state of California). Avoid contact with desirable foliage as injury may occur. Do not use on food or feed crops.

Broadcast: 1-2 pts. Solera Diquat Landscape & Aquatic Herbicide in a minimum of 15 gals. of water per acre. Add the labeled rate of a 75% or greater nonionic surfactant per 100 gals. of spray mixture. Use an adequate spray volume to insure

good coverage.

Method

[Broadcast](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[N.A.](#)