

SOYBEANS - HARVEST AID

General Information

USE INFORMATION

PARAQUAT CONCENTRATE is a liquid formulation containing 3 lbs. of active ingredient per gallon. It contains a nontoxic odor to help prevent accidental ingestions. It also contains an emetic (an agent which will induce vomiting if the product is swallowed).

APPLICATION

PARAQUAT CONCENTRATE is a contact herbicide for control or suppression of a broad spectrum of emerged weeds including most small annual broadleaf and grass weeds. It can also be used to suppress perennial weeds by destroying green foliage and as a desiccant/ defoliant at harvest.

Complete coverage of target weeds is necessary to get good control because PARAQUAT CONCENTRATE is a contact-type herbicide. It is also necessary to obtain complete coverage for good crop desiccation and defoliations. Undesirable weed control and undesirable crop desiccation/defoliation will result if improper application technique and/or application to large, stressed, or mown weeds are made. Refer to the following details for specific application instructions.

Thorough coverage of all green foliage is required for efficacious weed control and crop defoliation and desiccation because PARAQUAT CONCENTRATE requires actively growing green plant tissue to function. Drought-stressed weeds, weeds with little green foliage (i.e., mowed or cut weeds), or mature woody bark of trees and vines are unaffected by application with PARAQUAT CONCENTRATE.

There is no residual soil activity to affect later-planted crops or later germinating weeds because clay and organic matter rapidly tie up PARAQUAT CONCENTRATE.

ROTATIONAL CROPS

After the last application PARAQUAT CONCENTRATE, all rotational crops may be planted immediately.

RAINFASTNESS

Rain occurring 30 minutes or more after application will have no effect on the activity of PARAQUAT CONCENTRATE because it is rapidly absorbed by the weed foliage.

USE OF A NONIONIC SURFACTANT OR CROP OIL CONCENTRATE

The following should always be added and be used at the recommended rates or there will be a reduction in efficacy of PARAQUAT CONCENTRATE.

Nonionic Surfactant: Either add a nonionic surfactant cleared for the current use containing 50-74% surface-action agent at 0.25% v/v (2 pts./100 gals.), or add nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pt./100 gals.), of the finished spray volume for ground applications. Add a nonionic surfactant at 0.25% v/v (2 pts./100 gals.) of the finished spray volume for aerial applications.

Crop Oil Concentrate: For ground applications, add a nonphytotoxic crop oil concentrate cleared for the current use that contains 15-20% approved emulsifier, with 1.0% v/v (1 gal./100 gals.) of the finished spray volume. Add 1 pt. of crop oil concentrate per acre for aerial applications. For cotton harvest aid, do not use crop oil concentrate when using PARAQUAT CONCENTRATE.

APPLICATION TIMING

Applications should be made to small emerged weeds. Larger weeds more than 6 inches in height may be more difficult to control than weeds 1-6 inches in height. If possible, when green foliage is removed either from grazing or mowing, allow the weeds to grow 2-4 inches in height. Also, during harvesting forage or grain crops before spraying, weeds present in the field are also cut. Therefore, raise cutter bars as high as possible from the ground to cut stubble and weeds at a greater height, allowing sufficient green foliage to remain for applications.

BURNDOWN OF GRASS COVER CROPS OR VOLUNTEER CEREALS

The best results occur for control of grass cover crops or volunteer cereals when PARAQUAT CONCENTRATE is applied prior to tillering or after boot stage, especially with a wheat cover crop or volunteer wheat. Complete control may not be achieved with treatments made between tillering and boot stage. Complete control of perennial cover crops should not be expected.

ENVIRONMENTAL CONDITIONS

This product is active over a wide range of environmental conditions such as cool (below 55°F), cloudy or overcast weather. However these conditions will slow the activity of PARAQUAT CONCENTRATE.

SPOT SPRAYING

Refer to the following table if only small areas are to be sprayed with labeled applications.

Limitations, Restrictions, and Exceptions

SOYBEANS Harvest aid

Minimum Total Spray Per Acre

- Ground: 20 gals.
- Air: 5 gals.

Additional Precautions, Restrictions and Directions

- Indeterminant varieties: Applications should be made when at least 65% of the seed pods have reached a mature brown color or when seed moisture is 30% or less. Determinant varieties: Apply when plants are mature, i.e., beans are fully developed, 1/2 of leaves have dropped, and remaining leaves are yellowing.
- Injury will occur on immature soybeans.
- Mature cocklebur, especially drought-stressed plants, are tolerant to PARAQUAT CONCENTRATE and desiccation will not be complete. Always use the higher rate when treating cocklebur.
- Do not graze or harvest for forage or hay.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

15 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

.

Timings

[When at least 65% of the seed pods have reached a mature brown color or when seed moisture is 30% or less.](#)